



This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

### Usage guidelines

Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.

We also ask that you:

- + *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.
- + *Refrain from automated querying* Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.
- + *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.
- + *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

### About Google Book Search

Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at <http://books.google.com/>

QK

92

L3

UC-NRLF




B 4 599 187




BIBLIOTHÈQUE

ED. BONNET, D. M. P.


















Monsieur Pash.

hommage de l'Auteur.

GENERA

UNIV. OF  
CALIFORNIA

ET SPECIES PLANTARUM, QUAE AUT NOVAE SUNT,

AUT NONDUM RECTE COGNOSCUNTUR.

AUCTORE MARIANO LAGASCA.

---

*SUPERIORUM PERMISSU.*

MATRITI EX TYPOGRAPHIA REGIA.

ANNO MDCCCXVI.

To Mr. W. H. L. A. B. G. L. A. C.

QK92

L3

BIOLOGY  
LIBRARY

## LECTORI.

**E**n, Lector benevole, plantas fere omnes ab anno 1804 ad hunc usque diem in Regio Matritensi Horto nominatas, aliasque, aut prorsus ignotas, aut nondum ritè cognitatas, diagnosi certiori definitas, vel et contractis descriptionibus, synonymisque certis passim illustratas.

*Scabiosae* et *Knautiae* genera plantas fructificatione distinctissimas complectentia, proindeque certis limitibus omninò nullis circumscripta, juxta principia immortalis Vaillantio, atque neotericis Botanicis pro *Compositiflorarum* generibus distinguendis accepta; in *Lepicephalum*, *Asterocephalum*, *Scabiosam*, *Knautiam* et *Pterocephalum* genera dispescui: *Balmisam* ab *Aro* flore fructuque diversam, *Demetriam* ab *Astere* habitu fructificationeque distinctam, *Palaeoforiam* ab *Stevia*, *Alcinam* à *Wedelia* limitibus vel ab ipsamet natura impositis distincti: *Dyssodiae* genus reformavi, ejusdemque species in totidem fere genera immerito dispersas ad unum, idemque genus, jubente natura, retuli: *Steviam*, *Alcinam*, *Melampodium*..... characteribus certis (ni fallor) determinavi, speciebusque novis adauxi: synonyma aut nulla, aut certa attuli, generaque 26 prorsus nova descripsi, quorum notitiam manuscriptam, jam ab anno 1806, auditores nostri habebant.

Quamprimum à gravibus, quibus nunc distineor laboribus, tantillum sim vacuus; harum ipsarum plantarum, aliarumque in Regio Horto, in meis itineribus, atque in herbariis diu jam observatarum descriptiones pleniores in opere majori publici juris faciam. Tunc enim incorruptum sapientum Botanicorum judicium libenter amplectens, etymologias, correctiones, synonymaue adjungam, quae eorum observationes (quas mihi communicare velint oro, obsecroque) vel et librorum exterorum lectio, quorum nobis, propter bellum diutissimum, maxima est penuria, docebunt. Lectores siquidem monitos velim, me hac de causa maximam recentissimorum operum partem non vidisse, ut ex adjuncta breviationum explicatione facillè patebit.

Praeclarissimis Viris, qui ad Horti Matritensis restaurationem operam quamque benignè dederunt, gratias quas possum maximas ago; eosdem, caeterosque Europae totius fere Botanical et Agriculturae Professores, quibus litteras rarioraque semina missi, mittamque deinde, rogo obtestorque, ut eadem me indulgentia atque liberalitate in posterum prosequantur, illud donec, quod mihi in votis est, consequar,



nempe ut perfectionis fastigium pertingat Reg. Matr. Hortus, feliciter  
REGNANTE PRINCIPLE, cujus auspiciis Horti restauratio, et incrementa,  
Botanices et Agriculturae amplitudo, discipulorumque praemia opima  
debentur.

Vale, Lector benevole.

Dabam Matriti Kalendis Aprilis anno 1816.

## NOTARUM EXPLICATIO.

- All. — Fl. ped. — Allioni Flora pedemontana.  
 Anales. Anales de Ciencias naturales.  
 Anal. du Mus. de Par. — Annales du Musaeum de Paris.  
 Anthod. — Anthodium.  
 Asso. — Asso Synopsis stirpium indigenarum Aragoniae.  
 Barrel. Rar. Icon. — Barrelier Plantae per Galliam &c.  
 Bauh. Pin. — Bauhinus Pinax.  
 Boccon. Mus. di Piant. — Boccone Musaeo di Pianta.  
 Bompl. — Bompland et Humboldt Plantae aequinoctiales.  
 Boldo. Herb. — Balthasar Boldo Herbarium, in Regio Matritensi Horto asservatum.  
 Brot. — Felix Avellar Brotero Flora Lusitanica.  
       — Phytogr. — Phytographia.  
 Cabrera. — DD. Antonius Cabrera, Ecclesiae Gaditanae Canonicus Magistralis.  
 C. vel. Cav. Ic. — Cavanilles Icones plantarum.  
       — H. R. M. — Hortus Regius Matritensis, opus ineditum, centum constans tabulis cum descriptionibus.  
       — Herb. — Herbarium in Horto Regio Matritensi asservatum.  
       — Praelect. — Description de las plantas....  
 Cervantes. — Dominus Vincentius Cervantes, Horti Regii Mexicani Professor.  
 Cerv. sem. — Cervantes in seminibus ex Mexico missis.  
 Char. gen. differ. — Character generis differentialis.  
 Cl. vel Clem. — Simon de Roxas Clemente.  
 Clus. Hist. — Clussius Rariorum plantarum Historia.  
 Dodon. Pempt. — Dodonaeus Plantarum Pemptades.  
 D. — Dominus.  
 DC. vel Dec. vel Decand. Fl. fr. — Decandolle Flore française.  
       — Icon. Gal. rar. — Icones plantarum Galliae rariorum.  
 D. D. — Dominus Doctor.  
 Desfont. Atl. — Desfontaines Flora Atlantica.  
 Dill. Elth. — Dillenius Hortus Elthamensis.  
 Elench. H. R. M. — Elenchus Horti Regii Matritensis.  
 Ferr. Hesperid. — Ferrari Hesperides.  
       fig. — figura.  
 Flor. — Floret.  
 Fl. Per. vel Fl. P. — Flora Peruviana et Chilensis.  
 Forsk. Fl. Ar. — Forskhâhl Flora Aegyptiaco-Arabica.

Gaert. — Gaertner de Fructibus et seminibus plantarum.  
 G. N. — Genus novum.  
 H. vel Hab. — Habitat.  
 H. in N. H. — Habitat in Nova Hispania.  
 Hedw. — Hedwig.  
 Herb. Reg. Hor. Matr. — Herbarium Horti Regii Matritensis.  
 Horn. Hort. Haffn. — J. W. Hornemanni Hortus regius botanicus haffniensis.  
 Hort. Reg. Mexic. — Hortus Regius Mexicanus.  
 H. P. — Hortus Parisiensis.  
 H. R. M. — Hortus Regius Matritensis.  
 H. vind. — Hortus vindobonensis.  
 Hortul. — Hortulanorum.  
 Jacq. Fl. Austr. — Jacquin Flora Austriaca.  
     — Hort. Schoenbr. — Hortus Schoenbrunensis.  
     — Obs. — Observationes botanicae.  
 Juss. — Jussieu Genera plantarum.  
 Intr. — Introducta.  
 Lag. Varied. — Lagasca Variedades de Ciencias, Literatura y Artes.  
 Lag. et Rodr. — Lagasca et Rodriguez.  
 Lam. vel Lamk. Enc. — Lamarck Encyclopedie methodique.  
     — Ill. — Illustrationes generum.  
 Linn. sp. pl. — Linnaei Species plantarum.  
     — Hort. cliff. — Hortus cliffortianus.  
 Lobel. Icon. — Lobelius Icones.  
 Loeß. iter. — Loeffling Iter hispanicum.  
 Lud. vel Ludov. Née. — Ludovicus Néé.  
 M. Sessé. — Martinus Sessé.  
 Murr. — Murray, Sytema vegetabilium.  
 Observ. — Observatio.  
 Oed. Dan. — Oeder Flora danica.  
 Ord. nat. Juss. — Ordines naturales Jussieu.  
 Ort. Dec. — Ortega Decades.  
 Pal. Beauv. Agrost. — Palisot de Beauvois Essai d'une nouvelle Agrostographie.  
 Pers. Ench. pl. — Persoon Enchyridion botanicum.  
 Pluk. Phyt. — Plukenet Phytographia.  
 Pourr. — Pourret.  
 Recept. — Receptaculum.  
 Rod. — D. Josephus Rodriguez.  
 Sch. vel Scheuc. Agr. — Scheuchzer Agrostographia.  
 Scop. — Scopoli. Deliciae insubricae.  
 Sessé. — D. Martinus Sessé in seminibus missis.  
 Sw. vel Swartz. Synop. Filic. — Swartz Synopsis Filicum, Editio secunda.

**Tab. vel Tabul. — Tabula.**

**Vahl. Symb. — Vahl Symbolae botanicae.**

**Vaill. Act. Acad. Paris. — Vaillant Acta Academiae Scientiarum Parisiensis.**

**Var. — Varietas.**

**Vent. Hort. Cels. — Ventenat Choix des plantes du Jardin de Cels.**

**(VV.) — Vidi vivam.**

**(VS.) — Vidi siccam.**

**V. S. fl. et fr. — Vidi siccam floridam fructiferamque.**

**W. vel Willd sp. pl. — Willdenow Species plantarum.**

**Waldst. — Waldstein Plantae rariores Hungariae.**

**Ze. Ac. Matr. — Zea in Actis Academiae medicae matritensis.**

**⊙ — Planta annua.**

**♂ — Planta biennis: in caractere generico florem masculum indicat.**

**⌘ — Planta perennis.**


**♀ — In caractere generico florem faemineum significat.**

**2 — Arbor, Frutex vel suffrutex.**

**3 — Florem hermaphroditum indicat.**

**† Denotat plantam nondum apud nos floruisse.**





# GENERUM

## SPECIERUMQUE PLANTARUM NOVARUM

### AUT MINUS COGNITARUM

#### DIAGNOSES.

#### CLASSIS I. MONANDRIA.

##### ORD. MONOGYNIA.

1. *LOPEZIA minuta*: herbacea: foliis sparsis ovato-lanceolatis eroso-dentatis: floribus racemosis.

H. in N. H. ? ○. — Praedicto nomine acceptus ex H. P. ann. 1815.

2. *LOPEZIA oppositifolia*: herbacea: foliis ovatis dentato-serratis pierisque oppositis, superioribus sparsis; floribus racemosis.

L. annua. *Cer. Sem.* ann. 1814.

H. in N. H. ○.

3. *LOPEZIA miniata*: fruticosa: foliis denticulatis caulinis ovato-lanceolatis, ramulis lanceolatis. — *Petala miniata*.

Hab. in Nova Hisp. ? Semina missit D. Sessé. ann. 1804.

##### ORD. DIGYNIA.

4. *CORISPERMUM glomeratum*: glaberrimum laeve: foliis lineari-lanceolatis caule longioribus: spicis foliosis squarrosis caulem obtegentibus.

Patria ignoratur. Introducta ex H. P. an. 1806. *Corispermum satraloides* nomine. ○. (V. v.)

##### G. N. ASPICARPA.

*Character gen.*

*Corolla* (calix Lin. et Juss.) calyciformis, 3-fida, basi tribracteata. *Stamen* brevissimum hypogynum. *Anthera* solida. *Ovaria* duo, alterum constanter fatiscens. *Stylus*. O. *Stigma*.... *Nux* coriacea. monosperma.

— *Ord. nat.* ? Apocineae. *Juss.* ?

5. *ASPICARPA arens*. — *Planta* herbacea, perennis, volubilis. Caulis pilis urentibus hispidus. *Folia* opposita, breviter petiolata, cordato-ovata, integerrima, subtus praecipue pubescentia. *Pedunculi* axillares, 3-fidi, 3 flori, bracteati: *Stigma* neque nudis, neque armatis oculis unquam rimare potui, etai centies

perscrutaverim. *Semen* exalbuminosum, cuius integumenta duo corculos 2-3-5-septemve involvunt, haud secus ac in Citro.

H. in N. H. 11. — *Nux*. viperæ caput refert, unde Generis nomen.

#### CLASSIS II. DIANDRIA.

##### ORD. MONOGYNIA.

6. *GRATIOLA crassifolia*: foliis lanceolatis integerrimis crassis: pedunculis unifloris: caule erecto. — *Caulis* inferne radicans. *Stigma* bilamellatum.

H. locis udis circa Salmanticam. 11. — Floret aestate.

*SALVIA* † *Calicis* *dentibus* 5. *subaequalibus*. 7. *SALVIA Hispanorum*: albido-tomentosa: foliis lanceolatis crenulatis rugosissimis: calicibus cylindraceis acutis.

*Salvia cretica angustifolia*. *Cluss. Hist. r. pag.* 343. plantam nostram sat bene representat.

Abunde venit in collibus aridis mediocris altitudinis in Hispania. *Suffrutes*.

*Salvia*, *Salvia fina*, *Selvio* et *Madreselva*, vulgo ab Hispanis nuncupatur. — *Hujus plantae* campboriferæ infusum theiforme, vel et saturativum decoctum, optimum est remedium, quo nervi studiis nimis, curis, laboribus, vigiliisve protractis labefacti, optime, cito, nec injucunde, reficiuntur. Medici nostrates, quin et vulgus ipsum *Paralyssi* laborantibus folia *maritima* praescribunt.

† †. *Calycibus*, labio superiore 3-dentato.

8. *SALVIA odorata*: foliis ovato-oblongis subcordatis acuminatis erosius rugosissimis subtus niveo-tomentosis: caule superne paniculato.

S. odorata. *Hornem. Hort. bafn. v. i. pag.* 33. H. prope Bagdad. *Horn. loc. cit.* Introducta supradicto nomine ex H. P. ann. 1804. 11. — Caulis adscendens, superne paniculatus florifer. — Verticilli summi steriles.

9. *SALVIA parviflora*: foliis cordato-ovatis acutis erosius crenatis: bracteis verticillo

6-floro brevibus; corollis sublaetis calyce angustioribus.

Corolla minime viscosa, purpurea, labiis adproximatis, calyce paulo longior angustiorque. — Stamina longitudine faucis.

Ignoratur patria. 21.

† f. *Calicis labio superiore indiviso: Petioli basi subvix bicallosi.* (americanae.)

10. SALVIA *lineatifolia*: foliis cordato-ovatis acuminatis lineatis serratis, subtus tomentosis; pilis numerosis axillaribus terminalibusque densissimis.

Folia rugosa nonnumquam suborbiculata, subtus albicantia mollissima. Spicae numerosae breves, densissimae, pyramidatae, truncatae. Corolla pallide caerulea, minor quam in *S. polystachya Cav.*, cui nostra planta similis: galea et calix pubescens.

Introd. ann. 1814 ex seminibus a cel. D. *Vicentio Cervantes. S. eremiticae* nomine ex Mexico missis. 2.

11. SALVIA *melissodora*: foliis cordatis, ovatis obtusis rugosis subtus villosis; spicis terminalibus, verticillis multifloris tomentoso-viscidis subsecundis.

Intr. ann. 1804 ex seminibus a cl. DD. *Sesé et Mozinno* ex N. H. advectis. 5.

12. SALVIA *adglutinans*: foliis cordatis, ovato-lanceolatis acutis serratis subtus candidantibus; corollis pubescentibus, calyce inflato 4-plo longioribus; caule hirsuto-viscoso.

*S. amaethystina. Smirb?*

Folia, supra praecipue, villosa-viscosa, praesertim juniora. Verticillis subsexflori, bracteati. Bractae ovatae, acuminatae, concavae, calyce breviores, deciduae. Corolla coccinea fulgentissima. Stylus superne compressus, ibidemque utrimque barbatus. Stigma glabrum, galea longius.

H. in N. H. 5. — Introd. ann. 1804 ex seminibus a praeclaudatis DD. *Sesé et Mozinno* advectis. Ejusdem semina misit D. *Vinc. Cervantes* ann. 1814 *Salviae fruticosae* nomine.

13. SALVIA *lanceolata*: foliis lanceolatis remote obsoleteque crenatis reflexis; spicis, terminalibus secundis brevibus; bracteis lanceolatis pedicello longioribus; calycibus fructiferis superne compressis.

*S. lanceolata. Horn. Hort. bap. pag. 32?*

Caulis obtuse 4-gonus, 4-sulcatus, sulcis duobus oppositis latoribus, villos tenui pulverulento, uti rami, petioli et foliorum pagina inferior, vestitus. Folia fere conduplicata, caulina cito decidentia. Bractae adscendentes concavae. Calix fructifer inferne ventricosus. Corolla caerulea calyce paulo longior. — Ante florescentiam *S. pilosae* Cav. (*S. sideritis* W.) non assimilis.

Habitat in N. H. — Floruit in R. M. H. aestate anni. 1804. ex seminibus per D. *Sesé* missis.

14. MONARDA *citrifolia*: foliis rameis lanceolatis concavo-carinatis reflexis; floribus verticillatis; calycibus 5-aristatis fauce barbatis. — M. *citrifolia*. Cerv.

Planta herbacea anua, villos brevissimo pulverulento. — Folia primordially ovata, subcordata, obtusiuscula, late crenata, longe petiolata; caulina inferiora elliptica. Bractae numerosae. Corolla laete carnea.

H. in N. Hap. — Floruit vere in R. M. H. ex

seminibus per D. *Vicentium Cervantes*, praedicta nomine missis ann. 1814.

## ORD. DIGYNIA.

15. ANTHOXANTHUM *ovatum*: spica ovata debssa: vaginis glabris: foliis ciliatis:

H. circa Gades, ubi legit D. *Antonius Cabrera*. 6.

## CLASSIS III. TRIANDRIA.

### ORD. MONOGYNIA.

16. VALERIANA *rupicola*: floribus triandris: foliis inferioribus spathulatis integerrimis indivisis trifidisque: caulibus pinnatifidis integerrimis. *Lag. Varietates de Cienc. Literat. y Art. Ann. 2. núm. 22. pag. 112.*

Val. globulariaefolia. *Decand. Flor. fr. vol. 4. p. 236.* et secundum specimen missum.

H. ad saxorum rimas in cacumine montis *Peñafurada*, non procul ab *Aras*. — Floret Junio et Julio. 21.

17. LOEFLINGIA? *renifolia*: pentandra: foliis orbiculato-reniformibus.

H. in N. H. — Semina misit D. *Sesé*.  
18. CYPERUS *bulbosus*: culmo seminudo triquetro basi bulboso: umbella simplici: involucri triphylo longo: spiculis linearibus conglomeratis: squamis obtusiusculis. — Juxta *Cyp. Luzulae. Lin.*

α. capitulis pedunculatis erectis.

β. capitulis pedunculatis reflexis.

Patria ignoratur 6. Stylus 3-fidus.

19. CYPERUS *acutiusculus*: culmo triquetro superne nudo: umbella simplici: spiculis linearibus obtusis spicato-capitatis: capitulis pedunculatis reflexis.

Umbella 6-radiata: radis duobus infimis elongatis, recurvatis; 4 interioribus brevissimis. Spiculae circiter duodecimflorae. Involucrum diphyllum, umbella triplo fere longius.

H. in N. H. Semina misit D. *Sesé*.

## ORD. DIGYNIA.

20. PANICUM *scaberrimum*: panicula erecto-patente, ramulis trigonis strictis deorsum scaberrimis: glumis bifloris: radice tuberosa.

H. in N. H. — Introd. ann. 1804 ex seminibus per D. *Sesé* missis.

21. PASPAUM *pubescens*: spicis alternis secundis, basi pilis fasciculatis: floribus alternis geminis pubescentibus.

H. in Havana. 21. Introd. ann. 1804 ex seminibus per D. *Sesé et Mdr. Espinosa* missis. 21.

22. MILIUM *filiforme*: culmo adscendente radicante: vaginis hirsutis: panicula subdilatata patente filiformi: floribus breviter pedicellatis: calycibus ciliatis.

H. in N. H. 21. — Introd. ann. 1804 ex seminibus per D. *Sesé* missis.

23. MILIUM *microsperma*: culmo decumbente ramosissimo: panicula patula capillari: floribus muticis.

H. in N. H. 6. — Semina communicavit perill. D. *Sesé*.

24. AGROSTIS *microsperma*: culmo debili ramoso: panicula angustata: flosculis lanceolatis aristatis calyce ovato longioribus.

H. cum praecedenti. Vid. vivam ex seminibus per D. Serré datis.

23. *PHALARIS quadrivalvis*: glumis acutis integris: calice 4-divalvi; valvulis exterioribus duabus ovatis brevissimis glabris.

Habitat in agris, ad agrorum versuras, et juxta vias, circa Matritum, Murciam urbem, Orcellim, alibique in Hispania. — Gratum bellius atque pecoribus pabulum. Vulgo *Rabo de cordero* ab Orcellitanis dicitur.

26. *PHALARIS ciliata*: panicula spicaeformi cylindracea elongata: glumae valvulis lanceolatis aristatis, dorso ciliato-hispidis.

β. Gramen typhoides latifolium spica foliacea prolifera longissima Nob. *Scheuch Agr. p. 66*. Descriptio optima.

Patriam ignoramus. γ. — Gramen diversissimum à *Phal. pbleioides*. L.

27. *STIPA gigantea*: panicula nutante ramosa: aristis longissimis curvatis nudis: ligula obtusa extus tomentosa: foliis intus pubescentibus tandem convolutis.

*Stipa juncea*. Cav. *Præf. n. 89*. Venit in incultis circa Matritum, et praesertim in Praedio Regio la Real Casa del campo dicto. — Floret Majo et Junio. γ.

28. *STIPA capillata*: panicula patulo-erecta: aristis nudis rectis tandem curvatis subquadripollicaribus; ligula acuta glabra; foliis convolutis-filiformibus, intus pubescentibus. — Festuca juncea folio. *Seb. Agr. p. 151. tab. 3. fig. 13. A.* — *Stipa capillata* Linn. quoad descriptionem.

Hab. locis arenosis aridis circa Matritum. γ.

29. *STIPA pubescens*: foliis setaceis: aristis longissimis brevissime pubescentibus: pedicellis glabris scabris. — *Stipa juncea* Lin. var. *Desf. Fl. atl. 1. p. 98. tab. 28*.

Venit in collibus stecis, prope Matritum in el cerro Negro, et in Murciae Regno in las Anorias, Pretola, Chinchilla et Albacete oppidis: ibidemque *Stip. parviflora*, tortilis, et barbata. *Desf. pennata* et tenacissima. L. inveniantur. — Floret Maji et Junii mensibus. γ.

30. *ARISTIDA divaricata*: panicula divaricata incurvata, ramulis interne nudis: floribus adpressis alterne fasciculatis subsessilibus: aristis intermedia longiore.

H. in N. H. unde semina communicavit D. Serré.

31. *TRICHOPODIUM salmanticum*: panicula patente: glumis acuminatis subaequalibus, aristis calycina brevioribus.

H. locis arenosis humidis circa Salmanticam. γ.

32. *POLYPOGON panicum*: panicula subspicata cylindracea: glumis villosis, dorso aristatis: calycibus truncatis submuticis. — *Alopecurus panicus* Linn.

Venit abunde circa Matritum alibique in Hispania. γ.

33. *POLYPOGON elongatum*: panicula coarctata subspicata elongata: gluma glabra acuta indivisa: aristis calicinis glumacea vix dimidio brevioribus.

H. locis uliginosis juxta Albacete oppidum. — Floret Julio. γ.

34. *ANDROPOGON barbinode*: panicula oblonga contracta: pedunculis alterne geminis fasciculato-polystachyis: aristis contortis

culmo ramoso, geniculis dense barbatis.

H. in N. H. γ. — Semina communicavit D. Serré.

35. *ANDROPOGON laguroides*: panicula contracta, ramis simplicibus bifidisque fasciculato-polystachyis: pedicellis et spicularum rachyibus lanatis.

H. in N. H. γ. — Semina misit D. Serré. 36. *AIRA capillaris*: panicula capillari laxa trichotomo-ramosa subfastigiata: corollis muticis calyce carinato brevioribus. *Lagar. Varied. de Cienc. Liter. y Art. an. 2. n. 19. p. 29.*

β. culmo inferne radicante. Poa agrostidea. *Decand. Icon. Gall. rar. Fasc. 1. tab. 1.*

Habitat in montibus Manchae: β. locis uliginosis Galleciae. γ.

37. *AIRA refracta*: panicula nutante: corollis infra medium aristatis, aristis subaequali, basi pilosis: foliis radicalibus duplicatis refractis subpungentibus. *Lagar. loc. cit.*

Delectatur locis arenosis humidis, juxta Used et Torremocha oppida, et circa Legionem urbem. — Belluis pabulum gratissimum. — Floret vere. γ.

38. *AIRA lendigera*: panicula trichotomo-ramosa divaricata: corollis infra apicem aristatis, aristis longiore: foliis linearis setaceis laevibus. *Lag. loc. cit.*

Locis sabulosis humidis delectatur, prope Used oppidum in Aragoniae Regno, alibique in Hispania. γ.

39. *AIRA subtriflora*: panicula erecta confertiuscula: calycibus plerumque 3-floris: corollis basi pilosis infra medium aristatis; aristis recta floribus subaequali: foliis subulatis strictis. *Lag. loc. cit.*

Legi prope Torremocha in Castellia nová; non procul ab Burdongo oppido in montibus Legionensibus, et circa Legionem urbem. — A Majo ad Julium mensem floret. γ.

40. *POA mexicana*: panicula patentissima stricta: spiculis ovatis suboctofloris, oblongisque subquindecimfloris: flosculis trinerviis: foliis subtus laevibus; ligulis abbreviatis eroso ciliatis: culmo tereti ramoso.

*Antherae* violaceae: semina subelliptica, utriusque obtusissima, atropurpurea. *Valvulae* calycinae interiores persistentes.

Hab. in Imperio Mexicano, et tamquam spontanea crescit in R. M. H. γ. — Semina communicavit D. Serré an. 1804.

41. *POA pallida*: panicula rectangule patente pyramidata: spiculis linearibus subdecemfloris adpressis: rachis demum valvulis interioribus persistentibus pectinata. — Poa pectinata. *Pers. Encb. bot. 1. p. 91?*

Patriam ignoramus. γ.

42. *POA scariosa*: panicula elongata, contracta subsecunda: glumae valvulis scariosis cuspidatis: spiculis subseptemfloris.

Hab. circa Gades. γ. — Semina misit cl. D. D. Antonius Cabrera.

43. *POA sulcata*: panicula patula: spiculis suboctofloris: calycis valvula exteriori quinque-nervi: culmo vaginisque ancipitibus. — Gramen glaberrimum 3-4 pedale. *Poa sulcata* Pers. proximum.

Legi in sylvis tractus Valgrande dicti, non procul a Pajares oppido. γ.

44. *FESTUCA delicatula*: panicula secunda



spicata lanceolata: spiculis patentibus s-flo-  
ris aristatis: calycis gluma altera brevissi-  
ma: foliis lineari-setaceis, canaliculatis. *La-  
garo. Varied. de Cienc. ann. 2. num. 19. pag. 39.*

Fest. cynosuroidi. Desf. similis.

Venit juxta vias prope Guadarrama, Villa-  
castin et Arevalo oppida; tum et circa Ma-  
triturum, sed raro. — Floret Junio et Julio  
mensibus. ☉.

45. *FESTUCA Fenas*: panicula secunda: spi-  
culis oblongis sexfloris: floribus acuminatis mu-  
ticipis: foliis planis.

Hab. locis humidis et juxta rivulos Orce-  
llis, Murciae, alibique in Hisp. merid. — *Fen-  
as* vulgo. — Bonum pabulum equis et mulis.

46. *FESTUCA scabra*: panicula spicaefor-  
mi: spiculis teretibus undecimfloris aristis: for-  
mibus vaginis scabris.

H. in Imperio Mexicano. ☉. — Semina  
misit D. Sersé.

### KORYCARPUS, Zea.

*Char. gen. different.*

Flores paniculati. Gluma bivalvis multi-  
flora. Calyx bivalvis, valvula exteriore acu-  
minata. Stamina 2. Semen sacco coriaceo, api-  
ce rima hianthe arcuato vestitum.

47. *KORYC. arundinaceus. Ze. Ac. Matr.*  
1806. *Gramen* glaberrimum, perennans. Cul-  
mus teretiusculus, nodis subevanescentibus.  
Folia lanceolato-ensiformia, planiuscula,  
striata, intense viridia. Panicula racemosa,  
subsimplax, pauciflora; pedunculis in an-  
thesi erectis, seminum pondere uti panicula  
tandem inflexi. Gluma brevis, coriacea, uti et  
valvula exterior calycis. Petala (Nectarium  
L.) duo, ovata, obtusa, alba.

*Festuca diandra. Hornem. H. baffn. 1. p. 97.*  
Hab. in America.

Colitur subdico in R. M. H. — Ex semini-  
bus. 4 D. Touhin *Holcus sp. nov.* nomine  
missis an. 1803.

48. *BROMUS alopecuroides*: panicula cylin-  
dracea spicaeformi: glumis 4-8-floris: flos-  
culis dorso ciliatis vel asperis, breviter aris-  
tatis, supremo submutico.

Species meo sensu valde polymorpha. Hu-  
jus varietates sunt. *Festuca cristata*. Lin. et  
*Fest. piloides* Desf. quas ad Koeleriam retu-  
lit cel. Persoon.

Occurrit frequentissimum in Hispania. ☉.

49. *AVENA Cavanillerii*: panicula divari-  
cata: spiculis pendulis plerumque unifloris  
calycibus hirsutis aristatis; arista dorsali gla-  
bra glumam multoties superante. *Lag. loc. cit.*

Stipa arenaria. Brot. *Phytogr. n. 6.*

H. locis arenosis siccis veteris Castellae,  
Baeticae, Murciae Regni, et circa Matritum.  
Floret Junio. ☉. — *Spiculacae* plerumque uni-  
florae, nonnumquam 2-florae, bis tantum  
flosculis 3-6 stipatas vidimus in Reg. Matr.  
H. ann. 1814. et 1815. *Antherae* apice penici-  
llatae.

50. *AVENA birtula*: panicula subsecunda:  
glumis bifloris: calycibus hirsutis triaristatis,  
aristis duabus terminalibus setaceis.

Habitat locis rudertis, incultis, ad agro-  
rum versuras et juxta vias Matriti, Murciae,  
Orcellis, Gadibus, alibique. Floret primo ve-  
re. ☉. — *Gramen, Aven. strigosae* Linn. facie.

51. *AVENA altifolia*: panicula attenuata  
erecta: calycibus trifloris: flosculis aristatis  
basique pilosis, intermedio sterili; foliis  
convolutis filiformibus subpungentibus.

Venit ad saxorum calcareorum rimas in  
Urchillo monte, prope Orcelim, et in montium  
catena meridionali prope Murciam urbem,  
alibique in Hispania meridionali.

Vulgo *Laton* et *Laston* appellant Orceli-  
tani. Floret ab ultimo Februario ad Majum  
menssem. ☉.

52. *AVENA scabriuscula*: panicula con-  
tracta angusta: spiculis 3-floris: flosculis gla-  
bris: foliis margine involuto-subulatis sca-  
briusculis. *Lagas. loc. cit.*

Venit locis arenosis circa Legionem urbem.  
Floret Junio et Julio mensibus. ☉. — Flosculi  
apice bisetos.

53. *BROMUS pendulinus*: panicula subsecun-  
da pendula: spiculis lanceolatis compressis 3-  
5-floris; aristis subterminalibus.

Br. pendulinus. Sersé. — *Brom. compressus*  
*Elench. H. R. M. an. 1805.*

H. in Hispan. unde semina advexit  
D. Sersé ann. 1804.

### G. N. HYMENOTHECIUM.

*Char. gener. different.*

Flores terni, in racemum spicatum, secun-  
dum digesti, polygami. ☉ Intermedium, late-  
rales ♂, aut neutri. Omnibus Gluma 2-valvis,  
valvulis membraceis, calyce brevioribus,  
plerumque uniaristatis, uniflora. Corol. 2-val-  
vis, aristata, Stamina tria. Hermaphrod. Stylis  
duo: Stigmata pubescentia.

Locus, juxta Chrysimum. Pers.

54. *HYMEN. unisetum*: glumis calicibusque  
truncatis emarginatis uniaristatis.

H. in Imp. Mexicano. ☉. (V. S.) Semina  
misit D. Sersé.

55. *HYMEN. tenellum*: glumis acutis: caly-  
cis valvula exteriore uniarista. — *Cynosurus*  
*tenellus. Cav. H. R. M. 1. tab. 5. fig. 2.*

Hab. juxta Limam. ☉. (V. S.)

56. *HYMEN. trisetum*: calycis valvula ex-  
teriore, triaristata. — *Cynosurus gracilis.*  
*Cav. H. R. M. 1. tab. 5. fig. 3.*

Habit. in Mexic. Imperio. ☉. (V. S.)

57. *HYMEN. quinqueisetum*: calycis valvula  
exteriore 5-aristata.

Habitat in Mex. Imperio, ubi legit b. Lu-  
dovicus Nee. (V. S.) ☉.

58. *DIGITAR. foliosa*: spicis geminis gla-  
bris: floribus solitariis pedicellatis: culmo re-  
pente folioso; ligulis longe ciliatis.

Gluma bivalvis, valvulis acutis, aequalibus,  
tertia nulla. Folia glabra laevia, nec ad va-  
ginarum os pilosa. Pedicelli ad lentem villu-  
suli. Hinc diversa à D. *Distachya* et *paspaloi-*  
*de Pers.* quibus valde affinis.

Habitat in Havana, ubi legit. D. Balb. Bol-  
do. ☉. (V. S.)

59. *CHLORIS polystachya*: spicis pluribus,  
patentibus: calycibus flosculisque glabris, mu-  
ticipis: culmo compresso.

H. in N. H. unde semina misit D. Sersé.

60. *CHLOR. distichophylla*: spicis fascicu-  
latis plurimis; flosculis muticis: foliis inferio-  
ribus distiche imbricatis subsensiformibus.

H. in plantis Bonariensi et in Chilensi

**Regno. II.** — *Glumae valvula* exter. sub apice aristata. *Spicae* 24 et ultra subsupthameae. *Valv.* exter. flosc. hermaphr. ciliato-barbata.

61. **CHLOR. retusa:** spicis digitatis sub-senis: glumae valvulae exterioris retusa mucronata: flosculo hermaphrodito ciliato-barbato mutico.

Hab. in Bonariensi Regno. Ibi legit cum precedenti *D. Lud. Née*.

62. **CHLOR. dolichostachya:** spicis fasciculatis plurimis laxis longissimis: glumae valvulae exterioris aristata: flosculis longe aristatis, scabris.

Hab. in Phillip. insulis *D. Ludovicus Née* legit.

63. **CHLOR. orinita:** spicis fasciculatis plurimis: corollarum valvulae exterioris apice indiviso triaristato. *Lag. Varied. de Cienc. Liter. y Art. as. 2. num. 21. pag. 143.*

Hab. in Phillipicis insulis. *Q. D. Lud. Née* legit.

64. **CHLOR. foliosa:** culmo rigidulo vaginis foliorum numerosis vestito: spicis digitatis: subquinis longe aristatis.

Hab. in Porto-Rico. *II. Herb. Cavan.*

65. **CHLOR. rufescens:** spicis digitatis senis: glumis muticis: calicibus imberbibus: culmo compresso. *Lag. loc. cit.*

Hab. in Phylip. insulis.

66. **CHLOR. pubescens:** spicis fasciculatis plurimis: flosculi hermaphroditi valvulae exterioris dorso marginibus ciliata apiceque barbata: foliis pubescentibus. *Lag. loc. cit.*

Habit. in Peruvia. *Q.*

**BOUÏLOUA. Lag.**

*Character gener. differentialis.*

*Flores* spicati, unilaterales, polygami. *Gluma* 2-valvis, subtriflora. *Flosculi* hermaphroditi valvulae exterioris trifida, aut tridentata: masculi vel neutri bifida, 3-aristata: aristis duabus marginalibus, tertia ex emarginatura.

67. **BOUÏ. hirsuta:** glumae valvulae exterioris dorso tuberculis piliferis mucronato-hirsuta. *Lag. Varied. de Cienc. Liter. y Art. as. 2. num. 25. pag. 141.*

H. in Imp. Mexic. *Q.*

68. **BOUÏ. juncifolia:** culmo ascendente subtristachyo: spicis lanceolatis subpilosis: foliis convolutis filiformibus.

Habit. in Imperio Mex. *II.*

69. **BOUÏ. barbata:** glumis aristatis laevibus: flosculi hermaphroditi valvulae exterioris inferne ciliato-barbata. *Lag. loc. cit.*

Habit. in Imp. Mex. et Ins. Phillipicis. *Q.*

*Spicae* 2-3-4, alternae, lineares.

70. **BOUÏ. ovata:** culmo subpentastachyo: spicis ovatis: glumis extus pubescentibus.

Habit. in Imper. Mex. *II.* — *Culmi* basis tuberoso-incrasata.

71. **BOUÏ. simplex:** culmo erectiusculo monostachyo: spica terminali oblonga glaberrima. *Lag. loc. cit.*

Habit. in Peruvia. *Q.*

72. **BOUÏ. prostrata:** culmo ramosissimo prostrato: ramis monostachyis: spicis linearibus glaberrimis. *Lag. loc. cit.*

H. in Imp. Mexic. *Q.*

73. **BOUÏ. scorpioides:** culmo erecto filiformi monostachyo: spica lineari-oblonga spirally revoluta.

Habitat in N. Hisp. *II.*

74. **BOUÏ. litigiosa:** culmo subdiviso polystachyo: spicis linearibus patentibus: flosculo pedicellato rabdo longe triaristato. — *Aristida americana* L. Sw. Flosculi hermaphr. valvula exterior 3-dentata, dentibus aristatis.

Hab. in Jamaica *Swart.*

75. **BOUÏ. bromoides:** culmo polystachyo: spicis brevibus subovatis glabris, subcoctofloris: flosculo masculo longe triaristato.

Habitat in Acapulco in N. Hispania. *Q.*

76. **BOUÏ. racemosa:** spicis glabris racemosis pendulis. *Lag. loc. cit.* — *Chloris curtispindula*. *Pers. Enchyb. bot. 1. p. 87. Willd. Sp. pl. vol. 4. pag. 427.*

Hab. in N. Hisp., in Peruviae regno, Bonariensi planitie, et Phillipicis insulis. *II.*

## G. N. CABRERA.

*Char. gener.*

*Spicae* digitato-fasciculatae, secundiflorae. *Flores* singuli excavationibus racheos insculpti, sessiles. *Gluma* 1-flora 2-valvis: valvulis ellipticis, aequalibus, convexo-concavis, calycem paulo breviorum includentibus. *Cal.* 2-valvis glumae conformis. *Stamina* 3. *Styl.* 1. *Stigmata* penicilliformia.

*Obs. Genus DIGITARIAE Pers. affine, à quo floribus omnino sessilibus, atque racheos cavatibus duplici ordine antice insculptis distinguitur.*

77. **CABRERA chrysoblepharis.**

*Culmus* glaberrimus, laevis, inferne compressus, superne trigonus, multinodis. *Folia* lanceolato-linearia, acuta, utrinque striata, tuberculis piliferis dentato-ciliata. *Vaginae* carinatae, hiantes, superne praesertim tuberculato-hirsutae, suprema longior, glabra-laevis.

*Spicae* fasciculato-digitatae, 6-12; 3-4 pollicares, lineares, erectae. *Rachys* supra plana, subtus longitudinaliter trigona, atque omnino florifera, apice excepto: anguli tuberculis plurimis setiferis ciliati, cilia setacea, aurea; unde nomen triviale.

H. in America meridionali, ad Panamaldem et Aricam portum, ubi legit cel. *D. Lud. Née* (V. S.).

78. **DYNERA arabiqa:** spicis racemosis: spiculis 3-floris.

Habitat in Aegypto. *Q.*

In H. R. M. Vere floret.

79. **DYNERA Lima:** spica simpliciter: spiculis multifloris.

*Cynurus Lima*, Linn.

80. **ROTTBOLLIA cylindrica:** spica teretibusulata recta: glumis ovato-subulatis integris unifloris.

Hab. circa Gades, unde semina misit cl. *D. D. Antonius Cabrera.*

81. **LOLUM aristatum:** spiculis multifloris gluma univalvi duplo longioribus: flosculis ellipticis longe aristatis.

*Lol.* perenne Linn. *Q.* aristatum. *Willd. Sp. plant. 1. p. 462. — Vaill. Paris pag. 80. tab. 17. fig. 3. bona.*

Venit inter segetes in Gallogrecia, unde semina misit *D. Santamarina.*

*Tritici species sativae.*

*A. Semina valvulis callosis corticata. Rachys fragilis. Glumae cartilagineae.*

\* *Spica opposita compressa.* (1)

82. *TRITIC. monococcum*: gluma monosperma, valvulis bidentatis, dentibus rectis: semine utrinque compresso. — *Trit. monococcum.* Linn. (2)

Abunde colitur in Hispania. — *Espirilla, Carruon et Escaña* vulgo.

83. *TRITIC. Cienfuegos*: gluma disperma; valvulis subidentatis, dentibus unguiculatis, semine tumido.

Colitur in Asturum Principatu una cum *Tr. Spelta*.

84. *TRITIC. Baubini*: glumis truncatis dense imbricatis: semine tumido. — *Spica sericea. Spiculae* 4-5-florae, 2-3 spermae *Glumae* mucro brevissimus. *Rachys* eximie fragilis.

Colitur in Burgalensi Provincia?

\*\* *Spica parallela compressa.*

85. *TRITIC. Spelta*: glumis valde truncatis striatis distinctis: semine tumido. — *Trit. Spelta.* Linn. — Abunde colitur in Hispania septentrionali, et praesertim in Astur. Principatu. — *Escanda* vulgo.

B. *Semina matura valvulis denudata. Rachys flexilis.*

\* *Glumae coriaceae.*

86. *TRITIC. hybernum*: spica parallela compressa submutica: glumis gibbosis truncatis mucronatis basi coarctato-plicatis, exinde nervo longitudinali obtuse prominente percursis. — *Trit. hybernum.* Linn.

Abunde colitur in Hispania, praesertim in utraque Castella. — *Trigo chamorro, Pelón, Toleta &c.* vulgo dicitur.

87. *TRITIC. aestivum*: spica parallela compressa aristata: glumis gibbosis aristatis truncatis basi coarctato-plicatis, exinde nervo longitudinali obtuse prominente percursis. — *Trit. aestivum.* Linn.

Colitur eadem ferme copia, iisdemque in locis ac praecedens. — *Trigo candéal, Xexa, Membrilla, Barbilla, Perñan, Picbi et Pichon*, vulgo appellatur in diversis Hispaniae Provinciis.

*Spicae glabrae vel pubescentes.*

88. *TRITIC. Linnaeanum*: glumis ventricosis glabris brevibus parvis acutiusculis breviter mucronatis anguste carinatis.

*Spica* plerumque simplex, non raro ramosa (*Trit. compositum* Linn.) *Glumae* ovatae, à basi ad apicem vere carinatae, acutiusculae, nec truncatae, basi minime coarctato-plicatae, quibus notis à binis praecedentibus praecipue differt.

Colitur in Catalaunia praecipue, Aragónia, Castella veteri, et alibi. — *Redondillo, Rodonell, Pisana* vulgo. In hac, et sequenti, varietates spica composita *Racimales* et *Trigos del milagro* vulgo nuncupantur.

89. *TRITIC. turgidum*: glumis ventricosis pubescentibus brevibus parvis acutiusculis,

breviter mucronatis anguste carinatis. — *Trit. turgidum.* Linn.

Colitur in Catalaunia praecipue, Aragónia, Castella veteri et alibi. — *Redondillo, Rodonell* vulgo.

90. *TRITIC. fastuosum*: glumis utrinque compressis late carinatis mucronatis pubescentibus: spica longa subpyramidata: semine tumido.

Colitur in Hispania meridionali praecipue, *Fanfarron, Leucostachya Trebeles, Arisprietor, Arisnegros, Rispineros, Alonros, Salmerones, Cascavos, Finaños*. *Chrysostachya Trebeles* et *Rubionis* vulgo audiunt.

91. *TRITIC. Gaertnerianum*: glumis utrinque compressis late carinatis glaberrimis mucronatis: spica longa subpyramidata: semine tumido.

Colitur in Hispania, praecipue in Extremadura, Baetica, Murcia et Valentia. — *Leucostachya Blanquillos, Alagas*. *Cyanostachya Azulejos, Moratos et Xijonas*. *Chrysostachya Rojales* vulgo inter Hispanos audiunt.

92. *TRITIC. platystachyum*: spica opposita compressa confertissima abbreviata glabra: glumis utrinque compressis late carinatis.

Colitur in Regno Granatensi. *Chapado et Patianchuelo* vulgo.

Sequenti simillima: glaberrima, immo nitore quodam diversa.

93. *TRITIC. cochlear*: spica opposita compressa confertissima abbreviata pubescente: glumis utrinque compressis late carinatis.

Colitur in Granatensi Regno. *Trigo Cubareta et Patiancho* vulgo. *Spica* densissime imbricata, saepe concava, latitudine longitudinem fere aequante. — *Aristae* secundae.

94. *TRITIC. Cevallos*: spica pyramidata-teretiuscula longissima glaberrima: glumis utrinque compressis late carinatis bidentatis: semine longissimo.

Colitur in Regno Granatensi, et alibi per Baeticam. — *Trigo Moro* vulgo.

*Spica* omnium longissima, 5-7-pollicaris, laxiuscula, saepe subarcuata: rachis dentibus totis aequaliter longoque barbatis, barba à sesquilinea ad duas lineas longa. *Gluma* oblonga, calyce adjacenti r-a lineas brevior. *Semen* oblongum, durum, sex circiter lineas longum, sesquilineam circiter crassum.

95. *TRITIC. durum*: spica pyramidata-teretiuscula longissima pubescente: glumis utrinque compressis late carinatis bidentatis: semine longissimo. — *Trit. durum.* Desf.

Colitur in Regno Granatensi. *Trigo Moruno* vulgo.

Vix à praecedenti diversum si villositate demas.

\*\* *Glumae foliaceae.*

96. *TRITIC. polonicum.* Linn.

*Spica* subpyramidata, villosa. *Glumae*

(1) *Spicam parallela compressam* dicimus, cui ampliora latera racheos, ideoque spicularum plano parallela: cujus vero ampliora latera spicularum plano contraria glumas depressè imbricatas exhibent, *opposite compressa* nobis est.

(2) Certe: utrum villi praesentia characterem vere specificum semper et ubique ad Tritici species distinguendas praestet, nondum satis compertum est, etsi observationes à nobis hucusque institutae id suadere videantur. Dum vero experimentorum longior series rem in aperto ponit; an *TRIT. Spelta, monococcum* et *aestivum* L. in duas unumquodque sit species dispartendum, an *TRIT. Linnaeanum* à *T. turgido*, *fastuosum* à *Gaertneriano*, *platystachyum* à *cochlear*, *Cevallos* à *duro* in Naturae typo probe distincta necne, negotium alea plenum pronuntiamus.

compressae, oblongo-lanceolatae, angustae carinatae, 5-striatae, obtusae, brevissime mucronatae, circiter pollicares, duas lineas prope basim latae; altera valvula spiculum superante. *Rachis* dentes tribarbat; herba lineam longa. *Semen Trit. Cewalkor* similimum, sed paulo brevius.

Cultur in Legionensi Regno. *Trigo de Ponia* vulgo.

97. *TRITICUM spinulosum*: glumis aristatis 4-5-floris: flosculis inferioribus hermaphroditis aristatis: culmo frutescente.

*Triticum. Boldo herb.*

*Culmus* frutescens, rigidus, inferne vaginis foliorum distiche equitantibus vestitus. *Folia* rigida, plana, acuminato-subpungentia, denticulato-spinulosa, superiora edentula. — *Spica Trit. aestivi* similis, subpolicaris.

Hab. in Havana. cl. D. Balb. *Boldo* legit. 5.

G. N. CERVIA. Rodr. (1)

Ord. nat. Boraginaceae. Fuss.

*Char. gen. Calix* liber, 5-partitus. *Corolla* infundibuliformis, fauce nuda: Limbo quinquelebo, hypogyna. *Stamina* tria. *Styli* duo. *Capitula* bilocularia.

98. *CERVIA saturejaefolia*. Rodriguez.

Planta herbacea a spythama ad pedem alta. *Saturejae bortenris* facie.

H. circa Hispaniam O. (V. v.)

#### CLASS. IV. TETRANDIA.

##### ORD. I. MONOGYNIA.

99. *APHANES cornucopioides*: caule flexuoso erecto, stipulis amplexicaulibus subimbricato: foliis tripartitis ad stipulam sessilibus.

*Planta* annua, pilosa 1-3-pollicaris: — *Stipulae* cordatae, concavae, maximae, incisae, imbricatae caulem vestiunt. *Folia* tripartita, lacinialis 2-3 fidis, cum stipulis confluentia. *Floras* glomerati, axillares. *Semen* unicum. — H. in arvis circa Matritum. O.

100. *ANCISTRUM Acaena*: foliolis obovato-cuneatis sericeis serrato-incisiss, interioribus decrecendo brevioribus: spicis capitatis caulibus ascendentibus.

*Ancistr. Acaena. Elench. H. R. M. ann. 1803.*

*Obs. Acaenae trifidae. Fl. Per. similis.*

Hab. in N. H. 21.

101. *ANCISTRUM laevigatum*: foliolis obovatis lineatis inciso-serratis utrimque glabris laevibus: spica capitato-globosa.

*Ancistrum humile. Pers. Encypr. t. p. 30?*

*Stamina* 1-4, plerumque duo vel tria. *Antherae* atropurpureae.

Hab. in N. H. 21.

102. *PLANTAGO notata*: foliis linearibus remote dentatis: dentibus basi intus pilorum fasciculo stipitis: spicis ovatis lanatis.

*F. foliis linearibus angustissimis, subintegerrimis.*

*β. Major dentibus elongatis.*

Venit locis ruderalis et juxta vias circa Pulpi Pagum. à *Cuenas overa* oppido ad Heliocratam urbem eundo. O.

G. N. LEPICEPHALUS.

Ord. nat. Dipsaceae. Fuss.

*Char. gener. contractus.*

*Flores* capitati, supra receptaculum commune paleaceum: paleis arcte imbricatis, exterioribus sensim brevioribus, extimis plerumque vacuis, brevissimis, anthodii vices gerentibus. *Corollulae* 4-fidae. *Stamina* 4. *Semina* calyculo duplici obvoluta: exteriori exacte tetraedro, 8-dentato, dentibus alternis longioribus, plerumque aristatis: interiore supra semen coarctato, apice urceolato, multidentato.

*Char. gen. different.*

*Flores* capitati, supra recept. commune paleaceum, arcte imbricatum: paleis exterioribus sensim brevioribus, extimis brevissimis anthodii vices gerentibus. *Semina* exacte 4-edra, calyculo interiore apice urceolato multidentato coronata.

*Species.*

103. *LEPICEPHAL. atratus*: corollulis radiantibus: squamis ovatis villosis sphacelatis: foliis caulibus pinnatis: pinnis ovatis lanceolatisque inaequaliter serratis decurrentibus, scabiosa atrata. *Elench. H. R. M. an. 1805.* — Scabiosa elata. *Horn. bort. bass. 1. pag. 126.*

*Planta* herbacea orgyalis et ultra. *Folia* subtus praecipue hirsuta, imparipinnata: pinnulae 4 inferiores ovatae breviores, reliquae lato-lanceolatae, terminalis omnium major. *Flores* longissime pedunculati. *Squamae* ovatae, superne sphacelatae, interiores acuminatae, extimae obtusae non raro etiam floriferae. *Semina* hirsuta, calyculo exteriori dentibus octo, erectis, spinescentibus.

Admodum affinis *Scabiosae alpinae* L., à qua floribus radiantibus praecipue differt.

Habit. in Caucaso. 21.

104. *LEPICEPHAL. leucanthus*: squamis ovatis obtusis muticis: foliis subpinnatis, lacinis linearibus plerisque integerrimis — *Scabiosa leucantha*, Linn. — Habit. copiosae locis saxosis in Hispania meridionali, praesertim in monte S. Michaelis juxta Orcelem. 21. — *Flores* et Recept. ut in sequente.

105. *LEPICEPHAL. leucanthemus*: squamis ovatis obtusis muticis: foliis pinnato-lyratis: pinnis lanceolatis, inciso-serratis. — *Scabiosa leucantha*. *Decand. Fl. fr. 4. p. 225. Elench. H. R. M. pag. 14.*

*Recept.* squamae interiores ovatae, exteriores ovato-rotundatae, sericeae, extimae quatuor ellipticae, dorso viridi. *Capitula* praecedentis crassiora. *Corollulae* subaequales.

Hab. in Regno Valentino. 21.

106. *LEPICEPHAL. centaurides*: corollulis subradiantibus, laciniosis oblongis: foliis decurrenti-planatis, pinnulis lanceolatis integerrimis subtus pilosis. — *Scabiosa centaurioides*.

(1) Genus novum ex Boraginearum ordine naturalit, *Hidroleae* L. affine, à cel D. F. Josepho Rodriguez detectum, qui Cerviam dixit in memoriam clar. D. D. Josephi Cervi, Archiatris hispani, Academiae medicae allarumque Scientiarum Hispanensis fundatoris, atque Botanices promotoris.



des. *Decand. Flor. fr. 4. p. 224.* — Scabiosa transylvanica. *All. Pedem. n. 504. tab. 48.* huic pertinere mihi videtur, potius quam Sc. transylvanicae. L.

*Squamae* exteriores obtusae, interiores apice subulato-subaristatae. *Corolla* ochroleuca.

*Obs.* Admodum affinis scabiosae corniculatae *Waldest.* à qua corollis subradiantibus lacinii oblongis praecipue differt.

*Obs.* *Lepicephali* species sunt *Scabiosa alpina*, transylvanica, syriaca *Linn.* et corniculata *Waldest.*; eisdemque generi ex *Authorum* descriptionibus adjungendae mihi videntur *Scabiosa ustulata Thunb.*, Sc. rigida *Linn.*, attenuata. *Ait.* scabra *Linn. Suppl.* 2 humilis et decurrens. *Thunb.* 2 tartarica *Gmel.*, uraliensis *Murr.*, ciliata, et coriacea *Horn.*

ASTEROCEPHALUS. *Vaill.* Scabiosa L.

Ord. nat. Dipsaceae *Juss.*

Charact. gener. contractus.

*Anthodium* foliaceum, subduplici ordine 6-polyphyllum, aut monophyllum partitum vel fissum, tandem reflexum. *Corollulae* limbo 4-5-fido, plerumque inaequali. *Stamina* quatuor, raro quinque. *Receptaculum* commune paleaceum. *Semina* calyculo duplici obvoluta: exterior fauce foveolato-plicata; limbo ampliato, scarioso, denticulato, campanulato, aut hypocrateriformi: interior supra semen coarctatus, corollifer, limbo 5-7-aristato.

*Char. gener. differentialis.*

*Anthodium* simplex, aut subsimplex, lacinii aut foliolis foliaceis, tandem reflexis. *Recept.* paleaceum. *Semina* coronata calyculo exteriori limbo ampliato scarioso, et interiore 5-7-aristato.

*Species.*

\* Corollulis quadrifidis.

107. ASTEROCEPH. *diebotomus*: caule dichotomo: foliis caulinis subsessilibus subintegerrimis: capitulis alaribus subsessilibus, terminalibus pedunculatis.

a. *Scabiosa parviflora*: foliis inferioribus obovatis crenatis. *Desf. Fl. atl. 1. p. 219.* — *Scabiosa* dichotoma. *Elench. H. R. M. ann. 1803.* — *Horn. H. baffn. 1. p. 124.*

b. *Asterocephalus annuus* foliis imis senectutis retusis. *Vaill. Act. Acad. Paris. ann. 1722. p. 181.* — *Scabiosa alpina* hieraci folio. *Boccon Mus. di Plant. t. 120.* — *Var. a.* subpythamea. *var. b.* pedalis, vel parum amplius, caule inferne simplicissimo, foliisque inferioribus petiolatis, sinuatis, lyratisve: ramis plerumque trifidis, pedunculorum minoribus, oblongis, integris. Planta pubescens.

*Obs.* Huic primae divisioni adnumerandae *Scabiosa succisa*, et Sc. integrifolia *Linn.*

\*\* Corollulis 5-fidis radiantibus.

108. ASTEROCEPH. *pectinatus*: foliis carnosis pinnatifido-pectinatis canaliculatis: caule patulo. — *Scabiosa pectinata.* *Elench. H. R. M. ann. 1803. p. 33. et 1816. pag.* — Juxta *Asteroceph. divaricatum*, seu *Scab. sigulam* *Linn.* Hab. in Arabia. — Introducitur ann. 1802 ex seminibus à D. Thouin missis. (O.)

109. ASTEROCEPH. *pilosus*: corollae radio calycem superante: foliis pinnatifidis lineari-bus, summis linearibus indivisis: pedunculis longissimis cauleque pilosis.

*Folia* radicalia lineari-oblonga, obtusa,

superne utriusque dentata. *Caulis* inferiora subbipinnatifida. *Caulis*, pedunculi et foliorum nervi pilis longis, raris, basi tuberculo insidentibus obsidi. *Corolla* ex albo-caerulescens: radius magnus, lacinii tribus exterioribus amplis, rotundatis, inaequalibus, crenatis, undulatis. — *Scab. argentea* *Linn.* admodum affinis.

Hab. in Manchae Provincia. *Introd. ann. 1803 ex Hort. patavino Scabiosae albae* *Scop. nomine.*

110. ASTEROCEPH. *elegans*: corollae radio maximo: foliis integris, caulinis connatis lanceolatis: summis ovato-lanceolatis: pedunculis longissimis. — *Scabiosa caucasica.* *W. R. M. Elench. ann. 1805.* — *Scabiosa* connatis foliis omnibus integris, radicalibus petiolatis lanceolatis pilosis, caulinis remotis connatis subserratis, caule paucifloro, pedunculis longissimis, calycibus villosis. *Horn. H. baffn. 1. p. 128.* — Communicata ex Rusia ann. 1804.

111. ASTEROCEPH. *intermedius*: foliis subbipinnatis: pinnulis subtiliformibus caudiculatis: calycibus monophyllis 6-8 fidis: pappo 7-aristato subpiumoso.

*Scabiosa maritima.* *Elench. H. R. M. 1803.* Sc. tenuifolia *id.* 1805.

*Anthodium* monophyllum, corolla brevius. *Calyculus* exterior limbo scarioso extus pubescente, interior arietis 5-7-undique barbatus. — Planta *Scab. gramuntiae* L. facie, et quasi media *Pteroccephalus* inter et *Asterocephalus*.

Hab. in Boetica australiori. *W. (V.V.)*. *Introd. ann. 1804 ex seminibus missis per D. Petr. Gutierrez H. R. M. Solum. D. 7. Rodriguez ex Hispanensi Regno adsportavit ann. 1804.*

*Obs.* *Asterocephalo* adnumerandae omnes Scabiosae corollis 5-fidis praeditae à cel. *Pers. Ench. bot. 1. p. 180 et 181.* enumeratae, Sc. *pterocephala* et *papposa* exceptis: eisdemque generi adjungendae Sc. *canescens* *Waldest.* et Sc. *succulenta* *Decand.*

SCABIOSA. *Vaill.* Scabiosae spec. *Linn.*

Ord. nat. Dipsaceae. *Juss.*

Character gener. contractus.

*Anthodium* duplici, aut triplici serie octo-polyphyllum, foliolis alterne inaequalibus. *Corollulae* limbo 4-5-fido, inaequali. *Stamina* quatuor. *Receptaculum* hirsutum. *Semina* coronata calyculo proprio interiore, scyphi-formi, 8-multidentato.

*Charact. gen. differ.*

ANTHOD. subtriplici serie 8-polyphyllum. *Recept.* hirsutum. *Semina* coronata limbo calycis propter interioris scyphi-formi, 8-multidentato.

*Conspectus specierum.*

\* *Scabiosae* *Linn.* *Anthodio* hemisphaerico vel stellato-patente. *Coroll.* 4-fidis, radiantibus.

112. SCABI. *arvensis*: foliis pilosis pinnatifidis lucissimis. *Willd. sp. pl. Linn. pag. 550.* W. Hab. in arvis et pratis Hispaniae septentrionalis praecipue. (V. v.)

113. SCAB. *hybrida*: foliis villosis, inferioribus pinnatifido-lyratis dentato-serratis. — *Scab. hybrida.* *Allioni. Auct. p. 9.* — *Decand. Fl. franc. 4. p. 227.* O.

*Folia* superiora ovato-lanceolata subinte-

gerrima, acumine longo terminata, subintegra: ramea lanceolata, acutissima, integerrima, inferne ciliata. *Anthod.* hemisphaerium. 11-13-phyllum, corolla paulo longius. *Calculus* proprius exterior abbreviatus, pubescens, limbo denticulato (nec scarioso) ut in *Knautia orientalis* Linn., qua de causa primo ad *Knaut.* retulimus.

114. *SCAB. legionensis*: foliis lanceolatis longe acuminatis remote dentatis, inferioribus integris

*Anthod.* 18-phyllum, coroll. longius. *Caulis* brevissimus retroflexus hispidus. *Calculus* proprius interior dentibus octo, aristatis. *Folia* sequenti duplo et amplius angustiora. — Floridam iungi in sylvis montium legionensium, mense Junio. Eandem ex montibus Santanderensibus misit D. Barnaba Salcedo. § 11? (VV.)

115. *SCAB. rylvatica*: foliis omnibus indivisis ovato-oblongis deuto-serratis. *Willd.* loc. cit. — *Frag. Pl. aust.* 4. tab. 368. bona. 11. (VS.) Hab. in Navarrea sylvis.

116. *SCAB. ovatifolia*: foliis superioribus lato-ovatis acuminatis serratis sessilibus.

Præcedenti simillima, sed gracilior. *Folia* ovata (superiora paria duo tantum vidi) acuminata, serrata, basi acumineque integerrima, sessilibus: pedunculorum ovato-lanceolata, integerrima, latiora, breviora tamen et acumine longe breviora quam in *Sc. rylvatica* Linn. (VS.) Patrum ignoro.

117. *SCAB. longifolia*: foliis lanceolatis integris integerrimis. *Sc.* longifolia. *Pers. Ench. bot.* 1. p. 120. — *Wald. Pl. Hung. rar.* 1. p. 3. t. 5. — Caulis inferne glaber. *Anthod.* corolla paulo brevius, foliis pluribus lanceolatis, 3-plici serie positis. *Calculus* interior limbo hisuto, 10-setoso. *Waldst. loc. cit.*

118. *Knautia* Linn. Foliolis calycis erectis in tubum oblongum adproximatis.

119. *SCAB. orientalis*: corollulis 5-fidis quinque calyce longioribus: foliis pinnatifidis, integrisque. — *Knautia orientalis* Linn. (VV.)

120. *SCAB. propontica*: foliis superioribus lanceolatis integerrimis: corollulis quadrifidis denis calyci aequalibus. — *Scab. propontica* Linn.

#### KNAUTIA. Linn.

*Ord. nat. Dipsaceae. Fuss.*

*Char. gen. contractus.*

*Anthodium* duplici serie hexa-polyphyllum: foliolis alternis brevioribus, plerumque multifloris. *Corollulae* limbo 5-fido, inaequali. *Stamina* quatuor. *Semina* coronata calyculi interioris limbo in radios 8, pluresque elegantior plumosos soluto. *Receptaculum* villoso.

*Char. gener. differ.*

*Anthod.* duplici serie 6-polyphyllum. *Recept.* subnudum, villosum. *Semina* coronata calyculo interiore papposo plumoso.

121. *KNAUT. palaestina*: foliis lanceolatis integris. — *Kn. palaestina* Linn. sp. pl. VV. 1. p. 121. ②

122. *KNAUT. spatulata*: foliis integris rotundato-spatulatis tomentosis.

Habit. prope *Ellin* oppidum in Murciae Regno. In Regno Granatensi legit D. Simon de Roxas Clemente. (VS.) — *Subfruticulus* viz

palmaris, dense tomentosus, tomento candido. *Anthodium* 8-12-phyllum, corolla subsexdecimiflora, 3-plo fere brevius. *Corolla* radialis, fuscis 5-fidis, extus sericeis. *Pappus* coloratus, radiis 14.

123. *KNAUT. plumosa*: foliis superioribus profunde pinnatifidis. — *Knautia plumosa* Linn. — VV. sp. pl. loc. cit. 2

*Inquirendae.*

124. *SCABIOSA pteroccephala*: corollulis quinquefidis: caule procumbente fruticoso: foliis laciniatis hirsutis: pappo plumoso. Linn. Sp. pl. — W. sp. p. 1. p. 560.

*PTEROCEPHALUS. Vaill.* — *Scabiosae species* Linn.

*Ord. nat. Dipsaceae. Fuss.*

*Char. gen. contractus.*

*Anthodium* duplici serie 4-polyphyllum, multiflorum. *Corollulae* limbo 3-5-loba, plerumque inaequali. *Stamina* 3-4. *Receptaculum* commune paleaceum. *Semina* calyculo duplici obvoluta, anteriore abbreviato, 1-2-aristato, posteriori longis: interiore supra semina coarctato, in pappum multiradiatum, setaceo-plumosum soluto.

*Char. gen. different.*

*Anthodium* 2-plici serie 4-polyphyllum. *Receptac.* paleaceum. *Semina* aristata papposoque plumosa.

125. *PTEROC. diandrus*: corollulis subaequalibus, tri-quadrifidis diandris. — *Scabiosa diandra* foliis pinnatis, planulis subfiliformibus: floribus fuscis, corollulis 3-4-fidis subaequalibus diandris: seminibus unibariatis, pappo plumeis. *Lag. Varied. de Cienc. Ann.* 2. n. 19. pag. 39.

Habit. locis arenosis sterilibus, circa Martitum, et praesertim in la Real Casa del Campo et Valdelatas. Cel. D. Profes. Link ex Lusitania misit cl. Cavanillesio. ①

*Anthodium* 4-6-phyllum. *Corolla* caerulea. *Paleae* spinulatae, longe acuminatae. *Aristae* seminis 1-2, semipilicares. *Pappus* radiis 12-14, eleganter plumosis, longis.

126. *PTEROC. waitantii*: corollulis quinquefidis inaequalibus. — *Pteroccephalus annuus latifolius* Vaill. *Act. Paris. ann.* 1721. p. 183. — *Scabiosa papposa* Linn. sp. pl. — *Willd. loc. cit.* p. 560.

*Inquirendae.*

*Scabiosa pteroccephala* Linn. Vid. supra.

127. *GALIUM caucasicum*: caule scabro: foliis verticillato 6-9-nis linearibus margine revolutis scabris mucronatis: pedunculis umbellatis dichotomis.

Habitat in Caucaso. 11. — *Semina* ex Russia acceptimus ann. 1805.

128. *GALIUM capillare*: foliis inferioribus cauliculorumque sterilibus ovatis, 4-nis, superioribus 6-8-nis linearibus setaceis laevibus: seminibus villosis.

*Gal. capillare*. *Cav. Ic. vol.* 2. p. 73. tab. 191. f. 1.

*Gal. microcarpon* Vahl. *Symb.* 2. — *Gal. setaceum* Pers. *Enchyrt. bot.* 1. p. 128. 2.

Occurrit passim in monte St. Michaelis juxta Orcellim, alibique locis saxosis in Valentiae, Murciae, et Granatae Regnis. ①

#### BOLDOA Cavan.

*Ord. nat. Atriplices? Fuss.*

*Char. gen. differ.*

*Corolla* (Calyx Linn. et Juss.) tubulosa, tetragona, ore 4-dentato, persistens. *Stamina* 3-4, corolla longiora, altero latere anulo hypogyno inserta. *Stylus* filiformis longitudine staminum, indivisus, incurvus. *Stigma* simplex, acutum. *Utriculus*.

118. *BOLDOA purpurascens*: herbacea glabra: floribus glomeratis 3-4-andris. — *Cav. Hort. Reg. Matr. tab. 7.*

Hab. in Cuba Insula. Boido legit.

119. *BOLDOA ovatifolia*: fruticosa pubescens: foliis ovatis triplinerviis integerrimis: floribus 3-andris.

Hab. in N. H. ?

120. *BOLDOA lanceolata*: foliis lanceolatis subtomentosis: caule fruticoso.

Salpianthus arenarius. *Bompl. et Humb. Pl. aeg. 1. p. 155. tab. 44.*

Hab. in N. H. ubi legit D. Lud. Neé. ?

## CLASSIS V. PENTANDRIA.

### ORD. MONOGYNIA.

121. *ONOSMA tricerosperrum* strigosum: floribus cernuis calyce longioribus: seminibus tricoribus.

Vulgarissima planta juxta vias, circa Matritum, *Pedro Muñoz, Toboso, Albacete* opida, alibi in Nova Castella, atque in Murcia Regno. — Floret Junio et Julio. *Pl. Ojo de Lobo* vulgo.

122. *CYNOSLOSUS brassicaefolium*: glabrum: foliis imis spatulatis in petiolum attenuatis, reliquis cordatis amplexicaulis: racemis ebracteatis. — *Cynogl. lusitanicum. Vahl. Symb. 2. pag. 34* (non Linnaei).

Habitat in Regno Cordubense.

123. *CYNOSLOSUS valentinum*: foliis angusto-lanceolatis hirsutis, caulinis sessilibus: lacinis calycinis ovatis: spicis ebracteatis.

Planta herbacea *Anchusae angustifoliae*. Linn. facie.

*B. Cavanilles* legit in Regno Valentino. *Pl.*

124. *LITHOSPERMUM diffusum*: caule suffruticoso diffuso: foliis substrigosis linearibus basi latioribus margine revolutis: corollis calyce triplo longioribus.

Venit in dumetis prope *Aroas, Pajares*, et in perquam pluribus aliis Principatus Asturicensis plagis. — Floret a Junio ad Septembrem. ?

125. *ECHIMUM salmanticum*: foliis radicalibus lanceolatis: staminibus longissimis glabris: corollae fauce subinpervia. — *E. Echium lusitanicum* Linn. ?

Hab. circa Salmanticam. *Pl.* — Augusto floret.

126. *ECHIMUM paniculatum*: strigosum, superne paniculatum: ramulis alternis spica bifida terminatis: corollis calyce brevioribus: foliis summis lanceolatis subpetiolatis. — Venit prope Hispalim. *Pl.* (V. s.)

127. *ECHIMUM valentinum*: strigosum: foliis lanceolatis: corollis campanulato-infundibuliformibus subaequalibus calyce longioribus: staminibus corollam superantibus.

Clar. Cavanillesius legit in *Noveldae* et *Valldignae* arvis in Regno Valentino. — Floret vere. *Pl.*

128. *ECHIMUM molle*: sericeo-tomentosum: foliis lanceolatis acuminatis: spicis paniculatis: staminibus corolla longioribus.

Planta sexpedalis speciosa. *Broussonet herb. Cav.*

Habitat in *Palominos* prope *Taganana* in *Tenerife* insula.

129. *IPOMAEA turbinata*: foliis cordatis acuminatis integerrimis: caule pedunculisque subbifloris folio brevioribus muricato-aculeatis.

*Ip. muricata. Jacq. H. Sch. 3. tab. — Elenc. H. R. M. pag. 9.*

Caulis glaber, volubilis, aculeis recurvis muricatis, uti et pedunculi. *Pedunculi* 1-3-flori, inferne muricati, folio breviores, flori ferè erecti, fructiferi reflexi, superne valde incrassati, turbinati. — *Folia* subtus glaucescentia.

Affin. *Ip. bona nox*: à qua, foliis minime angulato-hastatis distinguitur.

140. *POLAEONIUM mexicanum*: foliis pinnatis polyphyllis, terminali trilobis: floribus cernuis: calycibus villosis-viscidis. — *Polaem. mexicanum. Cervant.*

Planta herbacea pubescens, pube viscida, subpedalis. — Semina praedicti nomine misit *cel. D. Vinc. Cervantes* Hort. Reg. Mexic. Professor. — Floruit Mayo et Junio an. 1815.

141. *POGONIA integerrima*: foliis lanceolatis integerrimis: pedunculis axillaribus confertis unifloris. — *Folia* coriacea, extorsum latiora, mucronata. — *Pedunculi* axillares, 2-3, conferti, uniflori.

H. in Nova Hollandia. ? (V. S. fl. et fr.).

142. *SOLANUM uniflorum*: pedunculis axillaribus unifloris: calycibus rotatis: foliis plicatis geminis subsessilibus ellipticis.

*Sol. uniflorum Serré.* — Planta herbacea, perennis, pilis raris brevibus scabrisculis praedita. — *Folia* nonnumquam solitaria. *Corolla* helvola, calyce triplo longior. *Barca* ovata, glaberrima, ovi gallinacei parvi magnitudine.

H. in N. H. — *Introduct.* ann. 1804 ex seminibus per *D. Serré* missis.

143. *SOLANUM Cervantesii*: caule fruticoso inermi: foliis ovato-lanceolatis utrinque acuminatis repandis: racemis extraxillaribus cymoso-umbellatis pedunculatis. — *Sol. microcarpon* Cerv. *Elencur H. R. M. 1803.*

Habit. in Nova Hispania: colitur subdio in R. M. H. — Floret toto fere anno. ?

144. *SOLANUM nicotianae-folium*: caule aculeato fruticoso: foliis oblique cordato-ovatis oblongis acutis inermibus subundulatis viscosissimis.

Habitat....

Colitur in R. M. H. — In Tepidario floret autumnò et hybere. ?

145. *SOLANUM viscosum*: villos-viscosum, aculeatum: foliis subbipinnatifido-sinuatis: pedunculis suboppositifoliis racemoso-multifloris: calycibus fructiferis subinfatis membranaceis. — *Sol. inflatum. Horn. H. b. 177. 1. pag. 221 ?*

Colitur in R. M. H. — Floret in Tepidario toto fere anno. ? — *Corolla* alba ampla.

146. *PHYSALIS foetidissima*: foliis geminis dentato repandis tomentoso-viscidis ovalibus: caule herbaceo superne paniculato.

Habitat in Nova Hispania. Colitur in R.

M. H. ubi florescunt. Veri. Semina misit prael. D. Sessé ann. 1803.

147. *PHYLLIS peruviflora*: ramosissima villosa: ramis dichotomis divaricato-reflexis: foliis ovatis subcordatis repando-dentatis. — *Phys. pruinosa* Linn. admodum affinis. — Habit. in N. Hisp. ☉

148. *CEANOTHUS caeruleus*: foliis oblongis subcordatis serratis, subtus tomentosis: racemis compositis pedunculatis. — Habit. in Nova Hispania. ☉. — Semina misit D. Sessé.

### CEVALLIA. Lag.

#### Character generis.

##### CALIX. O.

COROLLA (Linnæo, Jussieu aliisque calix diceretur) monopetala, supra germen coarctata, exinde decempartita: laciniis linearibus, planis, erectis, aequalibus, persistentibus.

STAMINA: Filamenta quinque, subulata, plana, aequalia, brevissima, laciniarum alterarum corollae basi interna inserta. Antherae conniventes, sagittatae, biloculares, longitudinaliter intus debiscentes, appendiculo membranaceo, cylindrico, cavo, terminatae.

PISTILUM: Germen adhaerens, ovatum, basi angustius, styli filiformis, filamentorum longitudine. Stigma capitatum, emarginato-subbilobum.

PERICARPIMUM: Nux à corolla formata, ovato-globosa.

SEMIN exalbuminosum, ovatum: Testa crustacea, nucleo accreta, obscure-viridis. Cotyledones duae, ovatae, extus convexae, intus planae, sulco longitudinali exaratae, carnosae, albae. Radicula supra brevissima obtusa (1).

#### Character differentialis.

GERMEN adhaerens. CAL. O. COROLLA 10-partita. Antherae appendiculo membranaceo, cavo, terminatae. Nux. sparsae, 1-locul., 1-sperma.

Syst. Linn. Class. Pentandr. Ord. Monogynia juxta Theodm.

Ordin. nat. Jussieu. i Boraginaceae!

149. *CEVALLIA sinuata*: strigosa: foliis alternis, subsessilibus, sinuatis, subtus tomentosis: pedunculis oppositifoliis, elongatis: floribus spicato-capitatis. Lag. *Varied* de Gien. nat. loc. cit.

Planta, ut ex specimine quod possideo conjectari potest, herbacea, ramosa, *Echinopsis* facie.

Ramus, quem describo, teretiusculus, villo brevi scaberrimo, setisque rigidis raris: siliis à taberulo calloso utrisque obtusis, ut et pedunculi, et foliorum pagina superior.

Folia alterna, subsessilia, oblongo-lanceolata, obtusa, sinuata, marginalibus revolutis, lineata, subtus tomentosa, tomento (in

siccis) in ferrugineum colorem vergente, sesquipollicem longa, vix pollicem dimidium lata.

Stipulae nullae.

Pedunculi oppositifoli, elongati, erecti, 3-5 pollices longi, rigidi, subfiliformes, apice in pedunculos 4-5 partiales involucrem mentientes, fissi. Pedunculi partiales, breves, superne bracteati, bracteis linearibus, strigosis, intus flores sessiles, plures, in capitulum globosum congestos sustinentes.

Corollae lacinae ciliatae, extus pilis canescentes, intus glabrae laeves.

Habitat in Nova Hispania, ubi legit. D. Lud. Nes. li?

Obs. Mirum certe videbitur me plantam germinem adhaerentem, proindeque staminibus perigynis stipatam, corollae orbatae ad *Boraginacearum* ordinem ammandare. Non adeo miratu dignum si fortasse putabunt, quid cum *Clas. Linnæo*, familiis, seu ordinibus naturalibus vix, ac ne vix quidem vel et à peritissimis Botanicis characteribus sufficienter definiri atque circumscribi posse autem. Id enim impedit, et impedit semper et ubique plantarum, omniumque entium naturalium affinitas ab utraque parte ut verbis clar. et immortalis Botanicorum Principis Linnæi loquar. *Universalis* clar. Juss. dictus character, seu habitus universalis consideratio affinitatem plantarum monstrat, et docum qualem naturalis ordine occupare debeant sat primo intuitu indicat: si vero fructificationis notae eandem ac habitus viam commonstrant, tunc *quaesitum* botanicam detectum esse putatur.

Nostra planta, prima facie *Echinopsis* generis speciem refert, at fructificationis characteres nedom ab hoc genere, verum et à Compositiflorarum et confinium ordinibus tam longe lateque separant. Neque *Elegans* associandum *Cevalliae* genus, ut fortasse aliqui sibi persuaderunt, nam habitus alique characteres repugnant, dum *Boraginacearum* ordinem amant.

Habitus et fructificationis notae, nempe plantae strigositas, et calice scissilis, ut in *Echio* vulgari: foliatio extipulata, simpliciter alterna et inflorescentia extraxillaris: antherae deinde sagittatae, appendiculatae, ut in plerisque Boraginaceis: Stigma capitatum, bilobum ut in non paucis ejusdem familiae generibus; atque ramis tandem structura exterior, ejusdemque interior fabrica — *Boraginacearum* ordini praecipue accommodata. At corollae absentia, germinis adhaesio, et staminum perigynia *Cevalliam* à Boraginaceis omnino excludere videntur: sed naturam audiamus, ejusque interpretem, *Methodi ordinum naturalium* Auctorem sapientissimum.

Corollae praesentia inter characteres classicos saepe saepius à clar. de Jussieu assumitur, hosque characteres *valentiores* esse debere, vel ab ipso operis illic (4) affirmat praefatus vir: qui tamen à natura mellora pos-

(1) Haec mea describendi ratio nonnullis forte minime ardebit: his ut corollae loco calycem legant, atque ubi de *perispermio* agitur sequentia. „Semen unicum, calyci accretum“ substituant praecor, et omnia omnibus, si fallor, plana fiant. Ego vero dum limites calycem et corollam inter non inveniantur, corollam cum clar. Cavanillesi dicam: Floris tegumentum organa pericarpia immediata cingens, cujusvis structurae et coloris sit.

(2) Genera plantarum. pag. XLII.

tea edoctus, dum de affinitate *Caryophyllacearum* cum *Amaranthis* disserit (1), aliisque in locis, corollae defectum plantarum affinitatem minime dirumpere pronuntiat; *Saxifragis*, *Caryophyllacis*, *Amarantibus*, aliisque non paucis sententiam confirmantibus. Germen superum aut inferum ad rem non facit, cum plures extent familiae utroque gaudentes. Staminum insertione totum iunilur difficultatis momentum. Plantas vero staminibus hypogynis et perigynis stipatas naturam coniunxisse contendimus, uti demonstrant *Saxifraga* et *Geum* Tournefortii, species utriusque in unicum *Saxifragae* genus à clar. Linnaeo associatae, quin ullus postea emunctioris naris botanicus eas segregare sit ausus; ad naturae nutum id factum fuisse calycis libertate ejusque in speciebus adhaesione varia plane evincuntibus. Sed eas liceat in duo genera dispescere, prout calyx inferus est, aut superus: nihil tamen minus utriusque generis affinitas arcta et inconcussa manebit. Loco natali et in Regio Matritensi Horto *Saxifragam Geum*, *stellarem*, et *cuneifoliam* Lionaei plures attente observavi, easque staminibus vere hypogynis toties praeditas vidi. Id, y aut mihi certissimum est, aut nihil de indaganda staminum insertione certum habeo. Exempla quam plurima alia eodem facientia, quae *Monocotyledones* (2) *Leguminosae*, alique *Dicotyledones* ordines suppediant, consulto ut praetermittam ipsa rei evidentia jubet. *Cenallium* itaque habitu, staminum, pistillorum, seminisque characteribus praecipuis, *Boraginis* strictè dictis omnino esse conformem neminem inficias iturum putamus.

150. *SALSOLA serifera*: floribus glomeratis digynis: foliis teretibus laevibus apice seriferis: axillis laniferis: caule laevi subangulato.

H. in calcareis aridis in Murciae et Valentiae regnis. C. *Barrilla fina* vulgo.

151. *SALSOLA tamariscifolia*: foliis sparsis subulatis de cideis: floribus axillaribus solitariis in spicas digestis. — Anabasis tamariscifolia. Linn. Sp. pl. — Cav. Ic. vol. 3. pag. 42. tab. 283.

Abunde venit inter saxa sabuloso calcarea, et juxta vias in australiori Hispania.

Floret primo vere. 5.

Vulgo *Erobilla* Murcitanis audit.

152. *ATRIPLEX polygama*: foliis linearibus, acutis repando-subdentatis, subtus canescentibus: floribus femineis glomeratis axillaribus. — Atriplex polygama *Sessé*.

H. in Nova Hispania. 5 Semina misit prael. D. *Sessé*.

153. *ATRIPLEX verticillata*: caule flexuoso herbaceo, ramis patentissimis: foliis deltoideo-oblongis eroso-dentatis subpilis postice integerrimis: calycibus faemineis, dorso margineque dentatis. — Atriplex verticillata. *Elemb. R. H. M.* 1803. — *Cav. H. R. M.* tab. 21.

Occurrit passim in Matriti suburbis et alibi in Hispania. C.

154. *HERNANDIA annua*: herbacea, annua, hirsuta: foliis ovatis ovalibusque subpetiolatis.

H. copiose in Hispania. C. *Diversa ab H. hirsuta*. L. et *Smith Fl. brit.* — Infusum urinae copiose ciet: areculas et calculos in vesica urinaria contentos eadem medela in quibus aegrotis expulsos vidi. Eundem ex ejus usu effectum obtinuisse perquam plures retulerunt inter quos, nominare sat est D. D. *Andram Ximenez* et D. D. *Emman. Berdè*, qui Orcelis medicam Artem summa cum laude profitentur. Vulgo *Quebrantapiedras*.

155. *CAMPANULA arvensis*: glaberrima: caulibus filiformibus: foliis cordato-rotundatis angulatis: floribus terminalibus pedunculatis solitariis: corollis patentibus capsulis 3-locularibus. *Lag. Varied. loc. cit.*

*Campanula cymbalariae* vel *haederacifolia*. *Pluk. Phyt. tab. 23. fig. 1.*

Venit ad saxorum rimas in el Puerto *Pajaros* et *Peñafurada*. Floret Julio. 21?

## ORD. DIGYNIA.

156. *LASERPITUM pubescens*: foliis ovatis sublancoelatis, basi oblique-cordatis serratis, subtus praecipue pubescentibus.

Hab. ad saxorum rimas in montibus *Legionensibus*. — Floret aestate. 21.

157. *THAPSIA tenuifolia*: glaberrima: foliis supradecompositis pinnato-multifidis, lacinulis canaliculatis plerisque tricuspidatis. — *Thapsia tenuifolia* lucide rigidoque folio hispanica. *Barril. obs. 639. tab. 156.* plantam nostram sat bene representat.

Habit. in Hispania meridionali, et praesertim juxta Sanctuarium de la *Fuen-Santa* prope Murciae urbem.

Floret Vere et aestate. 21.

158. *THAPSIA decussata*: foliis subbipinnatis, foliolis 1-2-pinnatifidis ad petiolum communem cruciatis lacinulis lanceolatis: petiolorum rachibusque hirsutis: involucrellis paucifoliis.

Habit. in Hispanici Regni. — Floret vere in R. M. H. 21. Semina adportabit cel. D. *Fos. Rodriguez*. Ad Fretum Herculeum legit D. *Ludov. Néb.*

## ANGELICA. Linn.

*Charact. gener.*

Fructus subrotundus. Semina quinque-angula: alae opacae, subfungosae, marginales latiores, dorsalis lateralesque primariae, marginalibus multo angustiores vel excissae.

*Obs.* Alae marginales latiores ex spatio intercostali laterali juxta costam marginalem primariam prodeunt, at tres reliquae ex costa dorsali lateraliisque primariis originem ducunt.

159. *ANGELICA villoria*: foliolis ovatis acutis acute serratis, subtus praecipue pedunculis pedicellisque villosis-scabris. — *Angelica sylvestris*. *Dodon. Pempt. pag. 314.* feon. bona. — *Angelica atropurpurea* Linn.?

Hab. juxta rivulos locis argillosis juxta *Pola de Lena* et *Pravia* oppida in Principatu *Asturicensi*.

(1) Id. pag. 80.

(2) *ECHINANDIA terniflora*. Ort. Stamina tria hypogyna, tria alterna perigyna profert, cujusmodi et alia ordinis ejusdem naturalis genera: vel rectius, stamina habent hypogyna, quorum tria inferne extusque calyci (Juss.) adhaerent.

160. *ANGEL. sylvestris*: foliis ovatis acuminatis distinctis acute serratis. — *Angel. sylvestris* *Linn. 16c. cit.* — *Smith. Fl. brit. 1. pag. 311.* — *Angelica sylvestris. Dodon. Pempt. p. 318.* Folia nimis brevia depinguntur, caeteroquin bona icon. & foliis incisio-serratis.

Hab. juxta rivulos cum praecedenti et in Pyrenaeis. *II.*

161. *ANGELICA major*: foliis ovato-lanceolatis argute inaequaliter serratis, extremo trilobato, inferioribus petiolatis: seminum alis subaequalibus. — *Angelica major. Dodon. Pempt. p. 318.* Icon bona. Alae latiusculae, marginales paulo latiores.

H. in Castella veteri: circa *Lumbreras* oppidum legit *cel. D. Joseph del Pozo.*

162. *ANGEL. Archangelica*: folio lobato impari lobato lateralibus subdecurrentibus ovato-lanceolatis basi cordatis. — *Ang. Archangelica Linn. Oed. Dan. tab. 206. bona. Smith. Fl. brit. 1. pag. 311. \*81.*

Habitat in Baetica. *Herb. Reg. Hor. Maur. 6.*

G. N. POZOA.

*Ord. nat. Juss. Umbelliferae. Pone Atrantiam.*

*Char. gener. different.*

*Fructus pyramidalis-tetragonus, calycis dentibus coronatus. Petala integra. Involucrum completum: subcoriaceum, ampum, dentatum, umbella simplicis longius.*

163. *POZOA coriacea*. — *Planta herbacea, subdondrantalis. Folia simplicia, cuneata, superne profunde dentata, longe petiolata coriacea, 5-nervi. Involucrum obsolete lobatum, crenato-dentatum, multinerve, nervis subramosis.*

Habitat in Andium tractu *Cordillera del Platanon* dicto. *II.*

CAPNOPHYLLUM. *Geertn.*

*Char. gener. diff.*

*Semina aptera, dorso diffuso, costis rugoso-scabris, valleculis sulcatis. Corolla flocculosa.*

164. *CAPNOPH. simplex*: caule simpliciter umbellis terminalibus. — *Capn. africanum. Geertn. s. 1. 85. f. 6.*

165. *CAPNOPH. dichotomum*: caule dichotomo-sulcato. — *Conium dichotomum. Desf. del. 1. p. 246. tab. 66.*

Habit. in Baetica. *○*

G. N. HAENSELERA.

*Ord. nat. Umbelliferae. Juss.*

*Char. gen.*

*Fructus scrotiformis, inflatus: semina costis capillaribus, valleculis planis.*

166. *HAENSELERA danaeiformis*. — *Habit. in montibus ditionis Lumbreras: oppidi Provinciae Nemaentinae. Cel. For. del. Pozo. legit.*

167. *ORENANTHE diffusa*: glaberrima caule diffuso, inferne planis: foliis caulicis pinnatis linearibus planis: umbella subnuda: umbellulis involucrentis: floribus subradatis: fructibus globosis.

Habitat in Baetica.

Colitur in R. M. H. ubi Junio feret. *II.*

168. *MIRRHIS odorata* foliis subtus villosulis: involucrellis lanceolato-subulatis: seminibus sulcatis. — *Scandix odorata. Linn. Sp. pl. WV. 1. p. 1449. — Jacq. Aust. V. App. p. 48. tab. 37 bona.*

*Myrrhis major, cicutaria odorata. Eaub. Pin. 160.*

*Myrrhis. Dod. Pempt. V. Lib. IV. cap. X. pag. 689.* — *Plantae habitum sat bene exprimit.*

Venit abunde juxta rivulos en *Roncorviller, Irati, Articus*. — *Floret Junio et Julio. Nees. (V. S.) 119.*

169. *MYRRHIS sulcata*: foliis subtus canotomentosis: involucrellis oblongis concavis setaceo-mucronatis: seminibus profunde sulcatis.

Habit. in montibus Cantabricis ad tractum. *Pefia mayor de Mena*, ubi legit D. B. Salcedo (V. S.).

ORD. TRIGYNIA.

170. *ALSINE molluginea*: caule patulo dichotomo: foliis linearibus crassis glaucis.

Habitat in Nova Hispania. In H. R. M. floret Vere. *○* Ex seminibus a cl. D. D. Sessé et J. M. Mocino allatis.

ORD. PENTAGYNIA.

171. *STATICE furfuracea*: caule paniculato ramosissimo villo brevissimo furfuraceo: foliis spatulatis retusis pubescentibus.

*Statice pruinosa. Fork. Fl. Arab. 1* Venit Alonae juxta vias, locisque ruderalis non procul à mare. — *Floret Autumno. II.*

CLASS. VI. HEXANDRIA.

ORD. MONOGYNIA.

172. *ALOE tuberculata*: foliis linearilimulatis acutis, acute cartilagineo-mucronatis inferne canaliculatis cartilagineo-tuberculosus. — *Corollae incurvae*. — *Affinis Aloe tuberculatae WV. sed notis enumeratis distincta.*

Ignoratur patria. *II.*

173. *ALLIUM palustre*: spatha monophylla submultiflora: scapo obtuse trigono foliis filiformibus subtrigonis brevior.

All. palustre. *Pourr.*

*Petala oblongo-lanceolata, acutissima, purpureo-cerulea. — Stam. simplicia, corollae breviora, inferne dilatata membranacea, monadelph.*

Hab. locis paludosis in Hispania septentrionali. — *Floruit Junio in R. M. H. ex seminibus à perili. D. Andrae Pourret missis.*

174. *NARCISSUS flavus*: spatha multiflora: nectario urceolato truncato integro petalis ovatis triplo brevior: foliis linearibus obtusis canaliculatis subtus striatis.

*Scapus teres, subcompressus, erectus, foliis brevior. Corolla flava. — Colitur in Mortis. II.*

175. *NARCISSUS Broussonetii*: spatha monophylla multiflora: scapo fistuloso ancipiti, foliis linearilanceolatis subaequali: nectario obsoleto.

H. in Africa unde semina misit cl. Broussonet. *II.*

176. *NARCISSUS juncifolius*: spatha uniflo-

ra: nectario campanulato lobato crispo petalis subaequali: foliis semiteretibus subulatis. Hab. in Baetica. 21.

177. *ORNITHOGALUM canaliculatum*: scapo nudo teretiusculo: racemo oblongo submultifloro: bracteis subulatis pedicello longioribus: foliis lineari-subulatis canaliculatis tandem involuto-junciformibus.

Ignorat patria.

Colitur subdio in R. M. H. ubi Aestate et autumnio floret. 21.

#### G. N. LAPIEDRA.

*Charact. gener. differ.*

Flores spathacei. Corolla regularis, hexapetaloidea, patens. Stam. erecta. Antherae sagittatae, incumbentes. Capsula adhaerens trilobata.

*Obs.* Corolla regulari stellatim patente: staminibus rectis, erectis, et antheris sagittato-bastatis ab *Amaryllide*, cui valde affine, abunde discriminatur: iisdemque notis a *Leucojo* satis distinguitur.

178. *LAPIED. Martinezi*. — Narcissus autumnalis albus latifolius. *Barrel. Icon.* 993.

Bulbus ovatus, tunicatus. Scapus compressus, anceps, glaberrimus, laevis, uti tota planta. Folia lineari-lingulata, scapo breviora, fascia longitudinali alba percursa. Spatha 2-3-phylla, suboctoflora. Pedicelli erecti, triquetri, spatha breviores. Corolla intus alba, patens, marcescens. Anther. luteae. Stylus et stigma album.

Hab. ad saxorum rimas subhumidas, in monte Arcis Saguntinae, prope Sanctuarium de la Fuen Santa juxta Algezars oppidum in Murciae Regno, atque non procul a Malacensi urbe legit acerrimus Naturae scrutator D. Felix Haenseler.

Floret Autumno. Colitur in R. M. H. 21.

179. *BERBERIS pinnata*: foliis pinnatis, spinoso-dentatis. — *Berb. pinnata. Elench. H. R. M.* 1803. pag. 6.

Habitat in N. H. 5. Circa Monterey et Nutka legit D. L. Nees.

### CLASS. VIII. OCTANDRIA.

#### ORD. MONOGYNIA.

180. *GAURA odorata*: foliis lineari-lanceolatis repando-dentatis: petalis quatuor ovatis adscendentibus.

*Gaura odorata. Serré.*

Hab. in N. H. 21. — Viget subdio in Reg. M. Horto. — Semina communicavit prae-laud. D. Serré ann. 1804.

#### ORD. TRIGYNIA.

181. *Polygonum serrulatum*: floribus octandris semitriginis: spicis subracemosis linearibus: foliis lanceolatis acutissimis serrulatis: stipulis vaginantibus setoso-barbatis.

Hab. in locis humidis atque inundatis, circa Murciam urbem, Orcellim, alibi in Hispania meridionali. — Autumnali et hyberno tempore viget. 21.

182. *CARDIOSPERMUM pubescens*: fruticosum, totum pubescens: capsulis obtusis.

*Corolla sanguinea, duplo major quam in Card. Halicacabo L.*

Hab. in N. H. — Semina misit D. Vincent. Cervantes ann. 1815.

### CLASS. IX. ENNEANDRIA.

#### ORD. TRIGYNIA.

##### G. N. ESPINOSA.

*Ord. nat. Polygoneae. Furr. Juxta Rheum;*

*Charact. gen. different.*

*Corolla* (calix *Furr.*) sexida; laciniis obovatis, aequalibus. *Stylis* tres, longi: *Stigmata* simplicia, acuta. *Semen*, 1. corticatum triquetrum.

183. *ESPINOSA verticillata*. — Planta herbacea, subdiorantalis, tomentosa. Caulis superne bifurcatus. Folia simplicia, subintegerrima, supra glabrescens; caulina superiora verticillato-terna, floralia quaterna. Florae terminales et in bifurcatione fasciculati: fasciculis involucri membranaceo obvolutis.

Hab. in N. H. 21. Cel. Lud. Nees legit.

### CLASS. X. DECANDRIA.

#### ORD. MONOGYNIA.

184. *CASSIA pubigera*: foliis quadrijugia subhirsutis, foliolis acuminatis ovatis, eximie lanceolatis: glandula subpedicellata conica ad petiolorum basim.

Hab. in Guayaquil. — (VV.) Semina communicavit D. Tafalla.

#### ORD. DIGYNIA.

185. *DIANTHUS caucasicus*: floribus solitariis: calycibus cylindricis acuminatis: squamis otonis ovatis acutis brevibus arcte imbricatis: petalis cuneato-oblongis, superne incisso-dentatis imbricatis.

Hab. in Caucaso, 21. — Semina accepimus ex Russia ann. 1805.

186. *ASCLEPIAS virgata*: glaberrima: caule erecto virgato: foliis oppositis linearibus sublanceolatis, margine revolutis.

Folia nonnumquam verticillato-terna. Umbellae laterales pedunculatae, multiflorae. Corollae albae, colore roseo-livido suffusae.

Hab. in Imper. Mexicano. Subdio viget in R. M. H. — Introd. ann. 1804. ex semin. a D. Serré datis. 5.

#### ORD. TRIGYNIA.

187. *SILENE geniculata*: caule basi subgynoso: foliis inferioribus sublinearibus ciliatis: spica secunda triflora: petalis bifidis.

*Sil. ciliata. S. geniculata. Dec. Fl. fr. v. 4. p. 756.* et secundum specimen a cel. Viro missum. — *Sil. arvensis. Lagasc. Varied. ann. 1. n. 22. p. 212.*

Die 2. Augusti mensis floridam fructiferamque legi in cacumine summi montis *Peñafurada*, non procul ab Arvas. 21.

188. *SILENE legionensis*: floribus spicato-secundis: petalis bipartitis: calycibus clavatis: foliis radicalibus lanceolatis acutis. cilia-



tis, caulinis lineari-subulatis. — Juxta *Sil. goudotiana*.

— *Caulis* perpendicularis; teres, gracilis, simplicissimus, brevissime pubescens, scabrusculus; subpedalis.

*Folia* radicalia numerosa in orbem prostrata, dense breviterque ciliata, in petiolum angustata; caulina paria tria dissita, connata, brevia, subulata; floralia subovata, breviter acuminata, brevissime ciliata, inferiora sesquilineam longa, superiora sensim breviora. — *Flores* 5-6 brevissime pedunculati, secundi, in anthesi patentes, postea erecti. *Calix* clavatus, subpollicaris, membranaceus 10-striatus, striis purpurascens, ad lentem vix pubescens, 5-dentatus, dentibus ovatis, brevibus, brevissime ciliatis. — *Planta* exsiccata tota flavescit.

*Hab.* locis incultis circa *Villadagor* oppidum Legionensis Provinciae. Floret Julio et Augusto mensibus. *ll.*

189. *SILENE hirsuta*: hirsuta: floribus spicato-secundis subsessilibus: calycibus ciliatis clavatis, fructiferis adpressis: petalis bifidis: corollae squamis cuilibet petalo binis. *Lagasc. Varied. loc. cit.*

Crescit in arenosis siccis juxta oppidum *Ghamartin*, in Hisp. meridionali. — In Lusitania.

Floret Majo. *○*.

190. *SILENE micropetala*: hirsuta: floribus subsessilibus spicato-secundis: calycibus ciliatis adpressis: petalis parvis profunde emarginatis. *Lagasc. Varied. loc. cit.*

*Silene micrantha*. *Linn. Can. Herb.*

Legi in tractu dicto *Barranco bondo* juxta viam, quae ducit ad tractum *Sacedal* non procul a *Fuencarral* oppido. — *Hab.* in Lusitania. *Herb. Cavan.*

Floret Junio. *○*.

191. *SILENE glauca*: floribus spicato-secundis pedicellatis: foliis lanceolatis acutis ciliatis: petalis bipartitis: calycibus floriferis cylindricis. — *Sil. glauca*. *Elecb. H. R. M. 1803.*

*Habit...* *○*. *Planta* glaberrima, laevis, glauca. — *Calyces* fructiferi clavati.

192. *SILENE longicaulis*: floribus racemoso-spicatis: calycibus ovatis glabris decem-striatis: petalis bipartitis acutis: foliis inferioribus spathulatis pubescentibus. — *Sil. longicaulis*. *Pourr. Elecb. H. R. M. ann. 1803.*

*Hab.* in Hispania. *○*.

193. *SILENE coarctata*: pilosa scabra: petalis bifidis: calycibus axillaribus subsessilibus hirsutis, fructiferis ovatis subglobosis, fauce coarctatis acuminatis erectis.

*Hab.* in Valentiae et Murciae regnis. *○*.

*Folia* inferiora lanceolata, acutula, in petiolum angustata; floralia lanceolata-linearia acuta calyce longiora. — *Calyx* fructifer ovato-subglobosus, fauce coarctatus, quinquefidus, lacinjis subulatis acuminatis, capsula longior, subcoriaceus, decemangulatus. — *Inter Sil. inclusam et Sil. cerastoidem* locanda.

194. *SILENE adscendens*: villosa-viscosa: foliis lineari-lanceolatis obtusis ciliatis: pedunculis unifloris fructiferis horizontalibus: calycibus oblongis ascendentibus: petalis bifidis.

*S. allophila*. *Clem. — Elecb. H. R. M. 1806.*

Peduncul fructiferi horizontaliter patentes. *Cal.* tubulosus, cylindraceus, basi circulariter reflexus, fructifer ascendens, oblongus, basi angustior. *Flores* hermaphroditi.

*P. S. de Roxas* Clemente legit in Baetica. *○*.

195. *SILENE divaricata*: piloso-glandulosa: caule divaricato dichotomo: petalis emarginatis: foliis ovato-lanceolatis acutis.

*Sil. divaricata*. *Clem. — Elecb. H. R. M. ann. 1806.*

*Sil. sedoides* *Juss.* affinis, sed duplo et amplius major quoad omnes partes. *Petala* rosea calycem superantia, lineari-cuneata, laeviter emarginata.

Venit locis arenosis in Baetica, unde semina adportavit. *cel. D. Clemente. ○*.

196. *SILENE macrophylla*: foliis cordatis pubescentibus latis: panicula dichotoma: petalis laciniosis: capsula depressis calyce inflato multo brevioribus.

*Viscago* fimbriata. *Hornem. Hort. bapn.*

*Hab.* in Caucaso. Floret Junio et Julio. *ll.* Semina acceptimus ex Russia ann. 1803.

197. *SILENE virida*: foliis lato-lanceolatis, acutis, supra nitidis: panicula trichotoma: calycibus piloso-viscidis: petalis bifidis.

*Sil. fruticosae* *L.* similima, à qua calycibus piloso-viscidis praecipue distinguitur.

*Hab...* Floret Majo mense in R. M. H. y.

198. *SILENE anastomosans*: glabra, caule paniculato virgato: foliis lanceolatis acutis, ramis linearibus: petalis bifidis cuneiformibus reticulato-venosis. — *Inter Sil. reticulatam et Sil. pitam* *Persoon.*

*Planta* herbacea, glaberrima, laevis, 1-3 pedalis, erecta. — *Calyces* clavati, acuti, purpureo-striati, superne striis reticulatis. — *Petala* bifida, cuneiformia obtusissima, laete rosea, venis purpureis eleganter anastomosantibus reticulata.

*Ex H. P.* acceptimus ann. 1801. *○*.

## ORD. TRIGYNIA.

199. *ARENARIA pungens*: foliis subulatis calycibusque mucronato-pungentibus: petalis calyce longioribus: stylis ternis quaternisque.

*Arenaria pungens*. *Clem. Elecb. H. R. M.* Habitat in summis montibus Regni Granatensis, praecipue in *Sierra nevada*. — Collitur in R. M. H. ex seminibus à *D. D. Simone* de *Roxas* Clemente missis.

Floret Vere et Aestate. *ll.*

## ORD. PENTAGYNIA.

200. *SPERGULA viscosa*: foliis verticillatis, villosa-viscosis: floribus decandris: petalis calyce longioribus. *Lagasc. Var. de Cien. an. 2. n. 22. p. 213.*

Legi Julio mense in locis arenosis, juxta nives in summitate montis vulgo de *Los pozos*, prope *Arcas*. *ll.*

201. *LYCHNIS dichotis*: petala emarginata: floribus solitariis dioicis: caule hirsuto: foliis radicalibus caulisque inferioribus spathulatis, summis lanceolatis.

*Hab.* in montibus Cerabi vicinis, locis depressioribus, in Valentiae Regno. à *ll.*

*Caulis* decumbens, hirsutus, florens ada-



cedens, superne dichotomus, subdodranta-  
lis. — *Folia* dense ciliata, subius praecipue  
subhirsuta, radicalia et caulinea inferiora spa-  
thulata, imperiolumat tenuata, superiora lan-  
ceolata, acuta, sessilia. — *Flores* peduncu-  
latis, erecti, solitarii, terminales et ex dichotomia. — *Corolla* rosea, ungulibus calyce lon-  
gioribus. — *Calyx* fructifer subglobosus.

## CLASS. XI. ICOSANDRIA.

## ORD. MONOGYNIA.

102. *Cactus humilis*: articulato-prolifer, humifusus: articulis obovatis oblongisque sublaemibus. — *Fructus* parvi, numerosi, obovati, purpurei.

Hab. .... 2.

103. *Inga leptophylla*: spinosa: foliis bipinnatis: partialibus 4-7-jugis, propriis 10-12-jugis: scytinis lanceolaris subfalcatis.

*Mimosa leptophylla*. Cav. H. R. M. tab.

38. — *Elench. H. R. M. 1893.*

Arborea subtripedalis in R. M. H., valde ramosa. *Spinae* stipulares binae, divergentes, recurvae. *Florum* capitula globosa, pedunculata. *Stamina* subquadrangula monadelphica. *Flores* omnes hermaphroditi.

Hab. in America. Semina misit cel. L.

Nec.

104. *ACACIA Siliquastrum*: foliis conjugatis 2-3-jugisque pinnatis: pinnis multijugis lineariibus obtusis: spinis stipularibus geminis rectis: spicis pedunculatis cylindraceis.

*Mimosa siliquastrum*. Cav. H. R. M. tab.

37. — *Elench. H. R. M. 1893.*

*Corolla* pentapetala. *Stamina* 10. *Pistilum* leviter pubescens. *Legumen* compressum, falcatum, polyspermum, glabrum.

H. in Chilensi Regno: viget subdio in R. M. H. ? — *Algarrobo de Chile* vulgo.

105. *ACACIA flexuosa*: foliis conjugato-pinnatis: pinnis suboctojugis, foliolis lineariibus obtusis basi angustioribus: spicis pedunculatis cylindraceis.

Arbor subtriflorialis, glaberrima, praecedenti similissima. *Spinae* stipulares, breviores. *Flores* decandri. *Legumen* teretiusculum, torulosum.

Habitat, coliturque cum praecedenti.

106. *ACACIA torquata*: foliis conjugatis bi-jugisve pinnatis, propriis multijugis lineariibus: spinis stipularibus binis: spicis cylindricis: scytino torquato.

*Mimosa torquata*. Cav. H. R. M. tab. 36.

*Petiolis* inter singula juga glandulosi, ultra folia protensi; uti et secundarii *Spinae* solitariae, breviter pedunculatae, decandrae. *Scytinum* in torquem convolutum, glabrum, externe inter semina sinuatum. *Semina* in singulo torulo solitaria, pulpa farinacea obducta. — An inde *Inga* adjungenda? Sed statim libera.

Habit. in America, unde semina adspostavit praed. D. L. Nec.

107. *ACACIA aculeaticarpa*: foliis bipinnatis: partialibus 2-12-jugis, propriis 6-12-jugis pubescentibus: leguminibus falcatis sparsim sutura utraque aculeatis.

Mim. aculeaticarpa. Ort. Decad. p. 134.

— Cav. H. R. M. tab. 39. — *Acacia aculeaticarpa*. Horn. H. bat. p. 449?

Hab. in N. H. ?

Obr. In hac, aliisque *Acaciae* speciebus, petala margine versus basin sibi inaequalem accumbunt, atque glutine, aut villo brevissimo, arcte connectuntur, ita ut corollam monopetalam mentiantur: at vi vel minima addibita, post anthesis praecipue, petala quinque absque ulla dilaceratione separantur.

108. *ACACIA penicillifera*: inermis: foliis bipinnatis, partialibus subduodecimjugis, propriis multijugis: petiolis eglandulosi subvillosi: dimarginatis: capitulis globosis pedunculatis polyandris. — Inter *Acac. desertorum* et *Ac. Hitchc. W. Sp. pl.*

— *Fruticulus* virgatus, debilis, subbipedalis. *Acac. Albicoides* W. facie. — *Folia* subius glaucescentia, utrinque glabra, 2-pinnata, pinnae primariae subduodecimjugae, secundariae 12-30-jugae, foliolis linearibus obtusis, ad lentem pilis raris ciliatis. *Petiolis* ad lentem, subvillosi, pilis adpressis.

Hab. in N. Hisp. ? ?

109. *LYTHRUM byssopifolium*: (Linn.) calycibus subsessilibus, fructiferis erectis teretibus, filiformibus capsula paululum brevioribus.

*Petala* lineari-oblonga, erecta, parva, 5-aut sex. *Stamina* quinque, sex, aut octo.

Hab. in locis udis circa *Albacete* oppidum.

*Murciae* Regni. — Floret Julio. ☉

110. *LYTHRUM flexuosum*: calycibus pedunculatis strictis, floriferis erectis, fructiferis horizontalibus. — *Rami* flexuosi. *Flores* dodecandri. *Corolla* hexa petala, rosea, petalis ovatis subcordatis.

Hab. cum praecedenti, eodemque tempore floret. ☉

111. *LYTHRUM acutangulum*: calycibus pedicellatis acute sexangularis, fructiferis infundibuliformibus capsula longioribus.

*Caulis* ramique acute angulati. *Folia* plerumque alterna, caulinea oblonga, rames lineari-lanceolata, acuta. *Flores* solitarii, breviter pedunculati, dodecandri. *Petala* rosea, oblongo-obovata.

Hab. cum binis praecedentibus, eodemque tempore floret. ☉

112. *CUPHEA bracteata*: caule retrosum hirsuto-patente: foliis ovatis acutis subundulatis: floribus solitariis interfoliaceis.

Hab. in N. H. Semina misit D. Sessé. ?

## ORD. POLYGNIA.

113. *POTENTILLA incisa*: tomentosa: foliis quinatis cuneiformibus jocularis, floralibus ternatis subsessilibus, triobis: petalis calycis subaequalibus: receptaculis hirsutis.

Habit. — Floret primo Vere in R. M. H. 2.

## CLASS. XII. POLYANDRIA.

## ORD. MONOGYNIA.

114. *Cistus glomeratus*: suffructicosus: capitulatus: foliis lanceolato-elipticis, subius praecipue incanis: floribus glomeratis axillaribus terminalibusque.

Habitat prope Acapulco et Cimapan in Nova Hispania. 2.

115. *Cistus sanguineus*: stipulatus herbaeae reclinator pubescenti-vilcidus: foliis omnibus oppositis: pedunculis axillaribus unifloris.

ris fructiferis retroflexis. *Lag. Variedades univ.*  
2. n. 19. pag. 40.

*Helianthemum retrofractum. Pers. En-*  
*chir.* 2. p. 78.

Venit locis arenosis siccis juxta oppidum  
*Cbamartin.* ○.

116. *CITRUS Calot*: petiolis alatis: foliis  
lanceolatis crenato-serratis acuminatis: fructu  
sphaerico parvo.

*Limon pusillus calaber. Ferr. Hesperid.*  
pag. 209. tab. rive pag. 211. Icon bona fructu  
majori minorique.

*Limon de carne*, et *Limon de confitar* vul-  
go dicitur.

Colitur in hortis orcelitanis.

Floret primo vere. 2.

#### ORD. TRIGYNIA.

117. *EUPHORBIA nutans*: caule patulo di-  
chotomo villosio: ramis apice floriferis nutan-  
tibus: foliis ovato-oblongis subcordatis obso-  
lete serratis: floribus axillaribus solitariis.

Habit. in N. H. ○.

118. *EUPHORBIA gemella*: caule prostrato-  
ramoso piloso: foliis oppositis oblique cor-  
dati inaequaliter dentatis: florum capitulis  
terminalibus geminis.

H. in N. H. ○.

119. *EUPHORB. pusilla*: umbella 2-3 fida:  
involucris oblongis obtusis, involucellis ova-  
tis: corollis 4-petalis: foliis cuneatis, superio-  
ribus angustioribus.

Habitat in Orcelitano monte Sti. Michael-  
is, juxta Seminarium conciliare ejusdem no-  
minis.

Floret Martio. ○.

120. *RESEDA lanceolata*: floribus 3-gynis:  
calycibus 7-partitis: foliis lanceolatis petio-  
latis: petiolis utrinque unidentatis.

Habitat locis montosis in Regno Jenensi  
et Granatensi. — Floret aestate. ○.

121. *RESED. erecta*: floribus pendulis tri-  
gynis: fructibus erectis oblongis: caule ram-  
isque virgatis: foliis tripartitis indivisisque:  
calycibus sexpartitis linearibus inaequalibus.  
Pedicelli setacei nutantes: fructiferi erec-  
ti, firmiores.

Venit in collibus cretaceis prope *Familla*  
oppidum in Murciae Regno, tum et circa  
*Aranjuez*, alibique in Hispania.

Floret Junio et Julio. ○.

122. *RESEDA media*: floribus 3-gynis: ca-  
lycibus corolla longioribus 6-partitis: petala  
superioribus capillaceo-multifidis: foliis  
integralis 3-lobisque.

Hab. in Hispania meridionali. ○.

#### ORD. POLYGYNIA.

123. *SEMPERVIVUM anomalum*: glaberrimum:  
foliis subulatis carnosis basi solutis:  
propaginis cylindricis: calycibus sex-decem-  
fidis.

Hab. in incultis sabulosis Regni Valentini,  
juxta *Cbamartin* et *Guadarrama* oppida, at-  
que in veteri Castella.

Floret Majo. 11.

#### G. N. BALMISA.

*Charact. gener. differ.*

*SPATHA* oblonga, tubulosa, apice cucul-

lato-fornicata: *Stam.* filamentosa, *antheris*  
reniformibus, unilocularibus. *Ovaria* stylifera.  
*Bacca* exsucca, membranacea, polysperma.  
*Spadix* imberbis.

124. *BALMIS vulgaris*: foliis cordato has-  
tatis: spathae apertura ovata: spadice incur-  
vo apice subclavato.

*Aris. serpentinum. Barrel. ic. tab. 573.* —

*Azum Arisarum. Linn.*

Hab. in arvis et locis ruderatis in Hispania  
meridionali.

Floret ab Autumno ad Ver usque. 11.

125. *CLEMATIS semitribas*: cithris opposi-  
tis scandens: foliis simplicibus cordatis serr-  
atis plerisque trilobis: ramis angulatis.

*Rami* angulati, atropurpurei. *Folia* pro-  
funde serrata. *Pedunculi* uniflori, axillares,  
solitarii, supra medium involucri, involu-  
cro bifido, lacinii rotundato-subcordatis,  
post anthesim latioribus reflexis. *Petala* 4,  
ovata, extus tomentosa. — *Clemat. cirriformis*  
Linn. similima.

Hab. 2 in Hispania meridionali? 2.

126. *CLEMATIS canaliculata*: suffruticosa  
scandens: foliis decompositis, foliolis linearibus  
canaliculatis: pedunculis axillaribus sub-  
trifloris.

H. in Murciae et Granatae Provincialis lo-  
cis saxosis, petrosisque. — Floret Vere.

#### CLASSIS XIII. DIDYNAMIA.

##### ORD. GYMNASPERMIA.

127. *TEUCRIUM subtriphyllum*: spicis oblon-  
gis: foliis oblongis tomentosis crenatis plani-  
nuculis oppositis, ternisque.

*Polium montanum luteum*, supinum ca-  
pite oblongo. *Barrel. obs. 320. Icon. 1082* bona.

*Teucrium trifoliatum Vahl?*

2. foliis verticillato ternis.

Habitat in montibus Jenensis Regni, ubi  
*Tomillo blanco* vulgo appellatur. — Floret Aes-  
tate et Autumno. 11.

128. *TEUCRIUM lanigerum*: caule capitu-  
lisque subovatis lanigeris: foliis crenato-plu-  
vatioidis linearibus oblongis revolutis.

Habit. in Regno Granatensi. *D. Clemente*  
legit. *Suffrutex* semipedalis, vel parum am-  
plius.

129. *TEUCRIUM compactum*: foliis cuneato-  
ovatis incisio-crenatis hirtis, floralibus  
petiolatis lanceolatis ciliatis: floribus spica-  
tis.

*Teucr. compactum. Clem.*

*Suffrutex* semi-1-pedalis, procumbens,  
pilis albis mollibus hirtis. *Folia* basi inte-  
gerrima, in petiolum brevem angustata: flo-  
ralla petiolata, lanceolata, acuta, ciliata,  
calycem superantia. *Calix* inflatus, fauce an-  
gustatus, semiquinquefidus; lacinii inaequa-  
libus, acutis, superiore ovata brevior.

Habit. in Granatensi Regno, ubi legit prae-  
laud. *D. D. Clemente.*

130. *SIDERITIS leucantha*: suffruticosa: ra-  
mis subcanis: foliis lanceolatis acutis mu-  
cronato-aristatis, utrinque 1-2-dentato-mu-  
cronatis: verticillis distantibus: bracteis sub-  
cordatis dentato-aristatis brevibus.

*Sld. leucantha. Cav. ic. 3. r. 304.*

*Spicae* elongatae flexiles, verticilli 6-flori.  
*Calix* subcampanulatus, aristatus. Corolla alb-

ba, galea semibifida, lacinialis divaricatis, obtusis.

Hab. locis aridis calcareis in *Urcillo* monte, alibique circa Orcelium.

237. *SIDERITIS angustifolia*: suffruticosa pubescens: foliis linearibus acutis integerrimis: bracteis cordato-rotundatis dentato-spinosis verticillo brevioribus.

*Sideritis montana* vermiculato folio. *Barrel. Rar. Icon.* 344. bona.

*Folia* inferiora conferta, breviora, obtusiuscula, subincana.

Habitat in montibus Regni Valentini, et praesertim circa *Canales* oppidum. — Floret Vere.

238. *SIDERITIS Tragoriganum*: suffruticosa subincana: foliis linearibus extrorsum latioribus subdentatis obtusiusculis mucronatis: bracteis cordatis serratis, dentibus aristatis.

*Sider. Tragoriganum. Elem. H. R. M.* ann. 1803.

α. Verticillis imbricatis.

β. Verticillis distinctis. — *Tragoriganum alterum. Cluss. Hist.* 1. pag. 355. — *Tragoriganum Cuslii. Lobel. Icon.* 494.

*Suffrutex* subpedalis, pube molli subincanus.

*Folia* caulinea duo supra lanceolata, serrata, qua nota facile a praecedenti et sequenti distinguitur. *Verticilli* numerosi, imbricati, aut distincti, adproximati, in spicam terminalem, 3-5 pollices longam digesti.

Habitat in Murciae et Valentiae Regnis. Circa *Tibi* oppidum legi, ubi *Rabet de Gat* vulgo dicitur.

239. *SIDERITIS linearifolia*: caule suffruticosa pubescente: foliis linearibus acutis mucronatis trinerviis integerrimis: dentibus bractearum calycumque spinulosis. — *Sider. linearifolia. Pers. Ench. bot.* 2. p. 117.

α. Verticillis dense imbricatis. — *Sider. montana hyssopifolia major. Barrel. rar. Ic.* 329.

β. Verticillis subdistinctis. — *Sideritis montana hyssopifolia minor. Barrel. rar. Ic.* 172.

*Caulis* pilis rariusculis pubescens, scabrusculus.

*Folia* virentia, subvittosa. *Bractea* cordato-ovatae, basi apiceque integerrimae, ad medium utrinque dentato-spinosae, calycibus longiores.

Habitat in montibus Regni Valentini, atque circa *Miranda de Ebro* oppidum alibique in veteri Castella.

240. *SIDERITIS foetens*: suffruticosa: foliis linearibus glabris acutis: spicis densis: calycibus frutescentibus fauce barbatis.

*Sider. foetens. Clem.*

*Suffrutex* sesquipedalis, caule ramisque virgatis, junceiformibus, glabris. *Pedunculi* villosi. *Spicae* cylindraceae, a sesquipollice ad quinque pollices longae, verticillis 12-24 et ultra dense imbricatis. *Bractea* ovatae, subcordatae, profunde incisae-serratae, dentibus subulato-aristatis, verticillo breviores. *Calicis* dentes subulato-aristati, faux pilis longis dense barbata. *Corolla* longitudine calicis, alba.

Prope *Nijar* oppidum legit. *D. D. Clemente*, in Granatensi Regno.

241. *SIDERITIS Cavanillesii*: subincana: foliis lanceolatis dentato-serratis: spica elon-

gata verticillis distinctis: bracteis cordato-orbiculatis spinoso-dentatis.

*Sid. scardioides. Cav. Ic.* 4. tab. 303. non Linnaei.

Habitat locis montosis Regni Valentini, circa Matritum, et in veteri Castella. ?

*Caulis* tomentosus, nec hirtus. *Folia* subtus praecipue tomentosa, magis minusve incana. *Corolla* flava.

242. *SIDERITIS spinosa*: foliis lanceolatis utrinque incisio-subidentato-spinosis: bracteis cordato-orbiculatis incisio-dentatis spinosis verticillo longioribus.

α. Caulis folisque tomentoso-hirsutis. — *Sider. spinosa. Vahl. Symb.* 1. p. 41. — *Lam. Dict.* 2. p. 168.

β. Caulis folisque glabrisculis. — *Sider. subspinosa. Cav. Ic.* 3. f. 109.

*Folia* extrorsum laetiora, ad medietatem dentibus mucronato-spinosis duobus tribusve utrinque excisa. *Spica* subpollicaris, verticillis imbricatis. ?

243. *SIDERITIS serrata*: foliis anguste lanceolatis utrinque acutis argute serratis, dentibus mucronatis: bracteis dentato-spinosis. — *Sider. spinosa. Cav. ber.*

*Planta* suffruticosa, hirsuta, exsiccata tota sordide flavet. — *Folia* lanceolata, angusta, utrinque acuta, apice mucronata, serrata, serraturis utrinque sex-octo, pluribusve, mucronatis, trinerviata, duplo longiora quam in praecedenti, tres lineas lata.

Habitat in montibus ditioribus *Tobarra* oppidi in Murciae Regno. ?

244. *STACHYS valentina*: foliis oblongo-lanceolatis, subcordatis hirtis: verticillis 6-7 floris summis sterilibus. *Lugas. Varied. loc. citat.*

Habit. in collibus *Enguerae* oppidi in Regno Valentino.

245. *SATUREJA obovata*: pedunculis axillaribus subternis secundis foliolosis: calycinis segmentis muticis: foliis carinatis obovatis obtusis integris.

Venit ad saxorum calcareorum rimas in australiori Hispania.

Floret Vere et Aestate. ?

*Saldorija* Orcelliani et Murciani appellant. Valentini *Herba d' oiver* vulgo audit. — Ad fructus oleae condendos adhibetur.

246. *THYMUS capitatus*: spicis capitatis bracteatibus: bracteis lanceolatis acutis ciliatis: foliis linearibus.

Juxta viam, quae ab urbe Alonensi ad *Melche* oppidum ducit legebam Majili mense anno 1810. ?

#### ORD. ANGYOSPERMIA.

247. *VERBENA polynemoides*: floribus axillaribus geminis sessilibus: foliis connatis subulatis subpungentibus.

*Suffrutex* ramosissimus *Polynemi arvensis*, facie. — Hab. in Planitie Bouariensi. (V. S.)

248. *VERBENA ligustrina*: foliis ovatis, oblongisque scabris margine revolutis integerrimis: spicis cylindraceis terminalibus.

Hab. in *Portu desiderato* ? (V. S.)

249. *VERBENA thymifolia*: spicis capitatis terminalibus siliatis: foliis ovatis ciliato-scabris margine revolutis: caule suffruticoso.

*Suffrutex* pygmaeus, ramosissimus *Sider-*

*niariae fruticosae* Lin. facie. — Habit. cum praecedenti. (V. S.)

244. *VERBENA tridactylites*: spicis capitatis terminalibus solitariis; foliis minutis connatis trifidis linearibus.

Fruticulus 1-2-pedalis, praecedenti, *Verb. thymi folia*: prima facie similissimus. — Hab. in insulis Maluins. (V. S.)

245. *VERBENA tridens*: spicis cylindricis, bracteis brevibus tridentatis; foliis trifidis laciniosis subulatis acerosis.

Frutex 3-4-pedalis. — Hab. in America meridionali. (V. S.)

246. *VERBENA juniperina*: spicis capitatis, bracteis indivisis calyci aequalibus; foliis tripartitis acerosis verticillum mentientibus.

Frutex erectus, ramosissimus, 3-6-pedalis. Bractaeae infimae duae, 3-fidae, reliquae indivisae. — Hab. in Audibus Chilensis, juxta viam, quae a *Cordillera del Portillo ad Mendoza* urbem ducit.

*Obs.* Omnes has Verb. species innumeras ex Herb. Cl. Cavanillesii accepti, lectas à cel. Peregrinatore D. Lud. Née. Earum fructificatio convenit omnino cum illa Verb. aff. *Pericarpium* omnibus est membrana utricularis, quae maturitate inferne evanescit, superne cum styli basi persistit seminibusque operculi adinstar superimposita, intusque impressionibus quatuor trigono-concavis insculpta est. 3 An rectius *capsula circumscissa* dicetur Verbenae pericarpium?

247. *LINEARIA filifolia*: foliis confertis linearibus filiformibus carnosius, inferioribus brevioribus quaternis, superioribus sparsis subulatis acicularis: floribus spicatis: calcare subulato rectiusculo longo.

Venit in collibus aridis circa Matritum. — Floret Majii et Junii mensibus. ☉

248. *MAURANDIA personata*: fauce clausa. — M. personata. *Serré*. — *Elencb. H. R. M.* ann. 1805.

Maur. antirrhiniflora. *Horn. Hort. buff.* 2. pag. 587.

Planta herbacea, glaberrima, laevis, scandens, minor quam *Maur. semperflorens*. Ort. quoad omnes partes. — Calycis laciniae lanceolatae, acutae, angustae, trinevartae, capsula longiores, corollae tubo breviores. Corolla caerulea, fauce clausa, triplo fere minor. *Maur. semperflorens*.

Habit. in N. Hisp. unde semina communicavit D. *Serré*, ann. 1804. — 21.

G. N. LAFUENTRA.

Ord. nat. Euphrasiae. *Fuss.*

Car. gen. diff.

Calix 5-partitus: corolla bilabiata, labio superiore bifido, inferiore trifido: capsula bilocularis polysperma.

249. *LAF. vorundifolia*. — Planta suffruticosa, villis densis apice glanduliferis viscidis angulique obsita, odorem amygdalarum amararum spirans, uni-bipedalis. Caulis decumbens. Rami assurgentes. Folia decussata, reniformia, crenata, crassa. Spicae axillares solitariae, breviter pedunculatae, oblongae, tantum elongatae, multiflorae.

Venit ad saxorum calcareorum rimas in monte Sancti Michaeis juxta Orcellim, et prope coenobium Sautae Catharinae, non

procul à Murcia urbe. — In Granatensi Regno legit D. *Clement*. Floret Autumno. 2.

250. *POTARRIUM agrimonifolium*: inermis: caulibus angulatis, pubescentibus: foliis argute dentatis, subius praecipue pubescentibus.

Pot. agrimonifolium. *Elencb. H. R. M.* ann. 1803. pag.

Habit. in Hispania? 21.

Pubescentia glandulifera. — Folia subius glauca.

251. *RANUNCULUS pubescens*: hirtus: pedunculis elongatis teretibus unifloris: calycibus pilosis reflexis: foliis tripartitis trilobis crenatis, summis lanceolatis integerrimis.

Habit. in Montibus legionensibus. 21.

## CLASSIS XIV. TETRADYNAMIA.

### ORD. SILICULOSA.

252. *ALYSSUM linearifolium*: caule suffruticoso patente: foliis incanis linearibus; siliculis ovalibus glabris.

Pl. suffruticosa pilis brevissimis stellatis incanescens. Caulis teres erectus, ramis patentissimis. Folia linearia, obtusa, extrorsum latiora, caudina latiora subrepanda. Siliculae ovales, obtusae, glabrae, laeves, 12-spermae, stylo brevissimo terminatae.

Habitat locis aridis prope *Moxente* oppidum Regni Valentini. *Cov. Herb.*

253. *IBERIS sempervivens*: frutescens: foliis linearibus acutulis villosis pulverulentis: integerrimis: pedunculo communi tereti: corymbo racemoso.

Iber. sempervivens. *Linn. Sp. pl.* — *Willd. Sp. pl. vol. 3. pag. 453.* Exclusis varietatibus et *Barrel. Synonymis*.

Caulis teres, cortice fusco-viridi, ramosissimus. Rami teretes, sparsi, numerosi, villos brevissimo pulverulento vestiti, uti folia pedunculi et pedicelli. — Folia conferta, linearia angusta, acutiuscula, integerrima, vix canaliculata, subius marginibusque scabriuscula, erecta. — Inflorescentia terminalis corymboso-racemosa. Pedunculus communis teres, nec sulcatus, multiflorus. Pedicelli compressi calyce longiores. — Corolla alba. — Calix corolla, stamina et stylus post anthesim purpurascens. Calix post anthesim persistit, non vero usque ad fructus maturatorem.

Habit. in locis montosis siccis circa *Aranjuez* oppidum, atibique in Hispania.

In Reg. M. Hort. floret à mensis Martii dimidio ad Majum usque. 2.

254. *IBERIS garrexiana*: frutescens, glaberrima: foliis linearibus subspathulatis integerrimis lucidis: pedunculo communi striato, angulato: corymbo racemoso.

Iberis garrexiana. *Allion Fl. ped. n. 920, tab. 40. fig. 3.* et *tab. 54. fig. 2.* — Thlaspi fruticosum sempervivens albo cratice flore. *Barr. ic. 734.* plantam juvenem primo anno florentem representat.

Habit. in Hispania. 2.

255. *IBERIS conferta*: foliis confertis linearibus lanceolatis acutis integerrimis denticulatisve eciliatis: pedunculis elongatis nudis: floribus corymbosis. *Lag. Varied. loc. cit.*

Habitat in dumetis prope *Arava*.

Floret Junio et Julio. 21.

156. *BISCUTELLA cuneata*: foliis cuneato-ovatis lanceolatisque subdentatis glabris: siliculis membranaceo-marginatis.

Herbacea, ramosa, tenera, glabra, laevis, annua.

Habitat in America meridionali. ☉.

#### ORD. SILIQUOSA.

#### G. N. DICRATIUM.

*Char. gen. different.*

*Corolla irregularis. Siliqua apice bicornis.*

157. DIC. *prostratum*: Planta herbacea, parvula, prostrata, pilis albis, adpressa scabrusculis undique obsita. *Folia* lanceolata, integra, extrorsum latiora, sparsa. *Spicae* solitariae, extraaxillares breves; primaria radicalis. *Calycis* foliola lutea sub apice fornicata ut in Herniaria. *Petala* lutea, exteriora duo calyce longiora submarginata. *Siliqua* adpressae, pedicello brevissimo, fere turbinato; compresso-tetragono sustentatae.

☿ *Erysimum bicornis. Willd. sp. pl. ?*

Venit locis ruderalis arenoso-calcareis, Orcellis juxta Coenobium Sancti Francisci, alibi; et Murciae in tuguriorum Lazareto del *Jilano de la Luz* dicto. D. D. Simon de Rojas Clemente eandem plantam in Granatensi Regno offendit. — Floret toto anno. ☉.

158. *SISYMBRIUM subastatum*: erectum glabrum: foliis caulibus runcinatis subintegerrimis: siliquis teretibus erecto-patulis: longissimis, stylo brevissimo globoso.

Patriam ignoramus.

Floret vere. ☉.

159. *SISYMBRIUM fugax*: glabrum laeve: foliis lyratis lobis lateralibus terminalique oblongo-lanceolatis acutis: siliquis filiformibus tandem contortis.

*Sisymbrium fugax. Elench. R. H. M. ann. 1805.*

Habitat in Hispania. 21.

160. *SISYMBRIUM rigidulum*: foliis pinnatifido-lyratis quinquelobis adpressis: siliquis rigidis patentibus brevissime pedunculatis: petalis calyce longioribus.

☿ *Sisymb. erysimoides. Desf. varietas ?*

Venit ad agrorum saepimenta locis siccis in *la Fuensanta* alibi; in Murciae Regno, atque circa *Cuevas overa* oppidum Granatensis Regni. ☉.

161. *HESPERIS ovata*: pubescens: foliis linearibus oblongis sinuato-pinnatifidis: siliquis pedunculatis subfiliformibus acutis.

Habit. in Baetica, locis sabulosis. ☉.

162. *HESPERIS repanda*: foliis canaliculatis repandis, inferioribus confertis linearilanceolatis dentatis, superioribus sparsis linearibus subintegerrimis.

Planta herbacea, perennis, subpedalis, fivida, pilis albis, brevibus, retroversis, adpressis scabruscula. — *Caulis* erectus, subangulatus, superne tetragonus. — *Florum* spica terminalis, multiflora, elongata; caule longior. *Calyc* atropurpureus. *Corolla* violacea petalis oblongis truncatis calyce longioribus.

Habit. in Montibus Marianis. 21.

163. *HESPERIS ornata*: incana ramosissima: foliis linearibus sessilibus, canaliculatis subintegris: siliquis filiformibus pedunculatis torulosis, mucronatis mucrone simpliciter.

Venit locis arenosis in tractu *Fuente Castellana*, alibi; circa Matritum.

Floret Majo et Junio. 21.

164. *SINAPIS dissecta*: siliquis ovatis torosis patulis cornu compresso-ensiformi brevioribus: foliis profunde pinnatifidis, lacinialis dentato-sinuatis pinnatifidisque.

☿. Caule siliquisve subhispidis pilis retrorsis.

Abunde venit inter sata, praesertim Lini usitatissimi, circa *Cuevas overa* oppidum, Helicratam, Murciam, Orcelem, alibi; Floret primo vere. ☉.

Vulgo *Morricanto* in *Cuevas overa*, oppido; *Rahuel* Orceletanis audit.

Herba plangae innuptae in acetaris adhibetur.

165. *SINAPIS heterophylla*: foliis lyratis hispidis, lobo terminali crenato-dentato, summis lanceolatis dentatis: siliquis filiformibus, pubescentibus adpressis.

Venit locis ruderalis circa Matritum. 21.

166. *SINAPIS subpinnatifida*: siliquis linearibus patulis rostro brevi obtuso seminfero: foliis lyrato-subpinnatis, lacinialis exterioribus sinuato-pinnatifidis.

Planta herbacea, annua, 1-2-pedalis, et ultra. *Caulis* teretiusculus, erectus, ramosus, pilis retrorsis hirtus. *Folia* lyrata, subpinnata, superne interrupte subpinnata, lacinialis longioribus oblongis, obtusis, sinuato-angulatis subpinnatifidisque, integerrimis vel dentatis, rachibus hirtis. — *Pedunculi* communes teretes, elongati glabri, laeves, racemosi-multiflori. — *Corolla* lutea. *Calyc* duplo longior. *Siliquae* lineares, tetragonae, nitidae, torulosae, 14-lineas longae, rostro obtuso seminfero, vix sesquilineam longo, terminatae. *Semina* minuta elliptica, rufa.

Hab. locis arenosis humidis umbrisque montium legionensium, praecipue circa *Murvas*. — D. B. *Salcedo* eam legit in Montibus Santanderensibus prope Mená. — Floret Julio. ☉.

167. *SINAPIS radialis*: scapo nudo: foliis radicalibus hispidis pinnatifidis: siliquis erectis laevibus cornu subulato longioribus.

Venit locis montosis juxta viam quae à *la Caragualda* ad *Chinchilla* urbem ducit in Murciae Regno, alibi; in Hispania meridionali. — Floret Junio. 21.

168. *BRASSICA laevigata*: foliis radicalibus, pinnatifido-runcinatis dentatis hispidis, caulinis linearibus subulatis, plerumque integerrimis: siliquis teretiusculis patentibus. ☉.

*Lagasc. Varied. loc. cit.*

Habitat in Castellà veteri, solo atreoso juxta *Medina del Campo* oppidum.

Collitur in H. R. M.

Floret Julio. ☉.

169. *Bruca erecta*: foliis radicalibus lyratis hirtis crenatis: caule erecto hispidulo: siliquis patentibus cornu monospermo duplo longioribus.

Venit ad agrorum margines subhumidos circa *Cuevas overa* oppidum.

Floret Februarii mense. ☉.

#### CL. XV. MONADELPHIA.

#### ORD. TRIANDRIA.

170. *SISYMBRIUM semisiliquosum*: radice;

tuberoso-fasciculata: caule superne ancipiti petalis planis lanceolatis.

Hab. in N. H. 21.

*Radix* tuberoso-fasciculata fibrosa; tuberculis fusiformibus, sordide albis; fibris coloribus, filiformibus. — *Caulis* gracilis, 4-8-pollicaris, superne anceps, simplex vel ramosus. *Folia* graminea, lineari-ensiformia, striatula, lineam lata, basi dilatata, vaginantis, caule longiora. *Spathae* 6-10, monophyllae, scariosae, oblongae, obtusae, in spicam oblongam imbricatam, uniflorae, exteriores duae carinatae, carina viridi, extrema acuta. *Corolla* lutea plana; petalis lanceolatis.

*Obs.* simile *Sicyr. micrantha* Cav. sed statura triplo majori, foliis planis, et radice tuberoso-fasciculata diversum.

#### ORD. PENTANDRIA.

271. *GOMPHRAENA decumbens*: caule decumbente brachiato, apice dichotomo: foliis ovatis: capsulis subglobosis sessilibus terminalibus sessilibus, dichotomaliibus pedunculatis.

*Gomphr. decumbens. Elench. H. R. M. ann. 1803. p. 17.*

*Caules* teretes, undique decumbentes, articulati, articulationibus incrassatis, rubicundi, pilis raris adpressis respersi, apice bifurcati. *Rami* consimiles brachiati. *Flores* capitati, capitulis ovatis, subglobosis, amethystinis, vel albis.

Hab. in N. H. 22.

#### ORD. HEPTANDRIA.

272. *PELARGONIUM inodorum*: umbellis capitatis multifloris: petalis calyci revoluti subaequalibus: foliis glabris cordato-orbiculatis crenatis sublobatis.

*Pelarg. inodorum. H. R. M. ann. 1803. pag. 27. — Hornem H. bot. 2. p. 634.*

Simillimum *Pelargonio althaeoides* W. a quo foliis glabris cordato-rotundatis obsolete lobatis praecipue distinguitur: folia enim *P. althaeoides* cordato-ovata, vel oblonga, sinuato-lobata, villosula sunt.

Habitat in Nov. Hollandia, unde semina adportavit D. L. Née. — Vere floret in hybernaculo. 23.

#### ORD. OCTANDRIA.

273. *LOURBEIRA cinerea*: foliis cordatis angulatis, trilobisque. *Ort. Decad. p. 108. Moenchae nomine.*

Hab. in N. H. 24.

#### ORD. DECANDRIA.

274. *ACALYPHA subtomentosa*: caule fruticoso subincano: foliis ovato-lanceolatis acumatis serratis subtomentosis: spicis solitariis axillaribus, supremis faemineis, inferioribus masculis.

*Acal. subtomentosa. Elench. H. R. M. ann. 1805.*

Hab. in Nova Hispania. 25. (Vv.) Semina misit D. Sessé.

275. *CAOTON patulum*: herbaceum albidum

tomentosum: foliis ovatis repandis basi subtus biglandulosi: fructibus porrectis cernuis.

Hab. in arvis in *Daimiel*, *Almagro*, et *Morab de Calatrava* oppidis Manchae Provinciae, circa *Algarres*, *Palmar* oppida in

Murciae Regno, alibi in Hispania.

Floret Septembri et Octobri. 26.

#### ORD. POLYANDRIA.

276. *ANODA punctata*: foliis inferioribus quinquelobo-hastatis, superioribus hastatis: petalis spatulatis: apice subdentatis, calyce subduplo longioribus.

*Corolla* punctata. — Diverissima ab *Anoda parviflora* Cav.

H. in Nova Hispania. — Semina misit praed. D. V. Cervantes.

277. *SIDA acutifolia*: hispida, foliis cordatis subpeltatis 3-lobis inaequaliter dentatis: pedunculis axillaribus: capsulis inflatis, basi birostratis. — *Corolla* caerulea.

*S. acutifolia. Elench. H. R. M. ann. 1805.*

Hab. in N. H. — Semina communicavit D. Sessé. 27. — In H. R. M. Vere, Aestate et Autumno floret.

278. *SIDA Serot*: foliis cordato-ovatis subtrilobis acuminatis dentatis subtus praecipue tomentosis: pedunculis axillaribus solitariis geminisque petiolo longioribus subbifloris. *Capsulis* quique.

Hab. in Nova Hispania. Collitur in R. M. H. ex seminibus a Perill. D. Sessé adportatis. Floret Autumno et hyberno tempore. 28.

279. *MAURA rubriflora*: pedunculis axillaribus brevissimis tri-multifloris: foliis cordatis angulato-subhastatis, crenatis: capsulis glabris inermibus.

Planta fruticosa orgyalis et ultra. — Affinis *M. Scopariae*, *scabrae*, *subhastatae* et *ovatae* Cav.: ab omnibus satis distincta, capsulis 8-9, glaberrimis, laevibus.

Hab. in Nova Hisp. Collitur in R. M. H. ex seminibus per D. Sessé adportatis. — Floret Vere. 29.

#### CLASS. XVI DIADELPHIA.

#### ORD. HEXANDRIA.

280. *FUMARIA parviflora*: calycibus minimis: floribus erectis: foliis pinnato-decompositis, foliolis tripartitis incis linearibus. *Fum. parviflora. Willd. sp. pl. 3. p. 868.*

Passim occurrit in arvis et ad agrorum versuras *Matriti*, *Orcelii*, *Alonae*, *Murciae*, alibi in Hispania. — Floret a Februarii mense ad Junium usque. 30.

281. *FUMARIA micrantha*: calycibus cordato-rotundatis corollae tubo latioribus: foliis pinnato-decompositis linearibus angustissimis.

*Flores* dense spicati, erecti. Venit in agris prope *Orcellum*, et circa *Murciam* urbem. — Floret primo Vere. 31.

282. *FUMARIA agraria*: calycibus oblongis acutis: pedicellis erecto-patulis bractea longioribus: foliis decompositis, foliolis cuneatis fissis.

*Flores* erecto-patuli, tandem recurvi. Quasi media inter *Fum. officinalem* et *capreaefoliam* L.

Frequentissime occurrit in agris circa Oracellum, Murciam. *Cucurbita ovata* oppidum, alibi in Hispania meridionali: ibidemque *P. officinalis* L. et *cucurbitata*. Linn. et Decand. frequentissime occurrunt. — Floret primo Vere. ☉.

## ORD. OCTANDRIA.

283. *POLYGALA linearis*: erecta glabra: floribus cristatis tribracteatis: racemis multifloris oppositifoliis: alis calycibus apiculatis uninerviis corolla longioribus: foliis linearibus obtusis.

Hab. locis montosis ab Anoriarum pago in Pretola oppidum eundo in Murciae Regno. Floret Junio. ☉.

## ORD. DECANDRIA.

284. *ERVUM tenuifolium*: pedunculis subtrifloris elongatis: leguminihus hexaspermis linearibus: foliis planatis linearibus: stipulis semisagittatis.

Ervum tenuifolium. Decand. herb. Venit inter segetes, locisque pratensibus Matriti, Oracelli, Murciae et alibi. — Ex Gallia misit cel. Decandolle. — Floret vere. ☉. Obr. a pecoribus bellisque non tangitur. Vulgo *Ojo de Sierpe* Oracellianis dicitur.

285. *LATHYRUS erectus*: pedunculis multifloris unifloris calyce brevioribus: cirrhis subulatis brevissimis dyphyllis: foliis, leguminibusque polyspermis linearibus.

Venit inter sata triticea circa Matritum. Floret Maio mensae. ☉.

286. *ONONIS vicifloris*: foliis pinnatis, foliolis oblongis obtusiusculis: stipulis bitidis subpetiolatis: pedunculis 1-7 floris: leguminibus lanceolatis glabris.

α. pedunculis unifloris breviter aristatis. *Vicia* calcarata. Desf. Fl. atl. 2. p. 166. — *Vicia monantha*. Retz. obs. 3. p. 39. — *Blanch.*

pl. Hort. R. M. ann. 1803. β. pedunculis 2-7 floris.

Venit inter segetes Matriti, Oracelli, in veteri Castellis, alibi frequentissima Alberja et Alberjana vulgo.

Floret a Februario ad Junium mensum. ☉. — Herba, Bobus praecipue, pabulum exoptatissimum: semina columbis escam praebent.

287. *PHASOLUS leptospermus*: caule tereti volubili ramosissimo hirsuto: foliolis deoidis integris obtusis: spicis axillaribus terminalibusque pedunculatis multifloris. — *Phas. microspermus*. Ser. d. — *Elench.* H. R. M. ann. 1805 et 1816, quod nomen retineri nequit, cum aliae plantae longe diversae a cl. Ortega antea inditum sit.

Hab. in N. H. unde semina misit. D. Sessé. (VV.)

288. *LUPINUS mexicanus*: calycibus alternis appendiculatis, labio superiore semibifido, inferiore obscure tridentato.

Lupinus mexicanus. Cerv. Juxta *Lup. Termis*. Linn.

Folia infima ternata, reliqua 5-7-nata: foliolis lanceolatis, mucronatis, supra glabris, subtus pilosulis. Stipulae setaceae, pilosae uti caulis. Pedunculus oppositifolius, superne racemoso-spicatus. Bractae deciduae se-

taceae. Flores alterni, breviter pedunculati, caerulei. Appendiculi setacei, breves. Legumina pilosa.

Habitat in N. Hisp. Semina misit praed. M. Vincent. Cervantes.

289. *CROTALARIA hirta*: caule erecto hirsuto: foliis ternatis, foliolis obovato-subrotundis: stipulis lineari-setaceis: racemis suboppositifoliis.

Hab. in Cuba insula Floridam fructiferamque habuimus in R. M. H. ab ann. 1805 ad 1808 ex seminibus a D. M. Espinosa missis. — Florebat autumnus. ☉.

290. *GENISTA terniflora*: racemis axillaribus subtrifloris: caulibus virgatis locanais: foliis simplicibus lanceolatis brevissime petiolatis: fructibus reflexis.

Suffrutex semibipedalis, erectus, *Anthyllidis cyttioideis* Linn. facile, sed omnibus partibus tenuior.

Hab. in montibus aridis in Valentiae et Murciae Regia.

Floret vere.

291. *ANTHYLLIS sericea*: fruticosa erecta: foliis planatis inaequalibus sericeis, floralibus digitatis: petioli spinescentibus: floribus capitatis.

Hab. in locis argillosis, cretaceis, juxta vias a Chinchilla ad Albacete oppidum eundo, alibi in Murciae Regno. ☉.

292. *ONONIS mollis*: pedunculis muticis unifloris longis: calycibus corollam aequantibus: foliis superioribus simplicibus, stipulis ovatis: fructibus cernuis.

Habit. in montibus saxosis calcareis in Murtae et Valentiae Regno.

Floret vere. ☉.

293. *ONONIS geminiflora*: villosa, foliis omnibus ternatis oblongis truncatis superne, dentato-serratis: pedunculis axillaribus bifloris aristatis petiolo longioribus: leguminibus oblongis induratis pubescentibus pedicellatis cernuis.

Hab. in locis aridis circa Matritum. — Floret primo Vere. ☉. — Flores ex albo-carnei, vexillum limbo roseis, eleganter striatum, dorso roseum. Carina acuta, apice rosea.

294. *ONONIS speciosa*: fruticosa: foliis ternatis ovatis duplicato-serratis: racemo subspicato terminali hirsuto: pedunculis subsolitaris bractea brevioribus.

Ononis. Haencler. — *Planta* fruticosa, pubescens. Folia ternata petiolata: foliola ovata vel elliptica, obtusissima, duplicato-serrata. Stipulae lanceolato-subulatae, basi vaginantes. Racemus terminalis, spithameus et ultra, multiflorus, bracteatus, hirsutus, hirsutiae glandulifero-viscosa. Bractae subcordatae, breviter acuminatae, concavae, pedicellis longioribus. Pedicelli inferiores solitarii, superiores geminati. Corolla lutea.

Habit. in Granatensi Regno, unde specimina sicca misit praed. D. Fel. Haencler.

295. *ASCHINOMENE hirta*: caule scabro hirsuto: racemis trifloris: bracteis serrato ciliatis: leguminibus hirsutis.

H. in Nova Hispania. ☉.

296. *ASCHINOMENE dentata*: floribus racemosis leguminum articulis rotundato-ellipticis muricatis: foliis pinnatis, foliolis obovato-cuneatis incisio-dentatis subtus punctatis. *Planta* herbacea. Caulis, rami, petioli,

pedunculis et pedicelli villosi, villo brevissimo apice glandulifero. *Folia* impari-pinnata, foliolis, 9, obovato-cuneatis, truncatis, superne inciso-dentatis, subserratis, junioribus subvillosis. *Racemi* terminales, multiflori, erecti. *Legumina* articulis 8-12, elliptico-rotundatis, punctato-muricatis.

Habitat in Amer. meridionali. 20?

397. *AESCHYMONENE hispidula*: pedunculis axillaribus unifloris: leguminum articulis orbiculatis crinitis: foliis pinnatis subcristatis: foliolis oblongis apice subdentatis. — *Petiolus* communis apice cirrho brevissimo simplicissimo.

Habit. in Amer. meridionali.

*Planta* herbacea diffusa. *Caulis*, petioli et pedunculi spalis raris, brevibus, retroversis hispiduli. *Folia* pinnata: pinnulis 8-12 oppositis oblongis, obtusis, mucronulatis, integerrimis, aut apice parce denticulatis: nervo marginibusque pilosis. *Petiolus* communis apice cirrho brevi, simplicissimo terminatus. *Legumen* lineare, articulis subsenis, orbiculatis, medio setis subplumosis stipatis.

Habit. in America meridionali.

398. *AESCHYMONENE papposa*: racemis terminalibus glabris multifloris: leguminum articulis truncatis setosis, setis plumosis.

*Planta* herbacea, perennis. *Radix* simplex, descendens, planta longior. *Cauliculi* villosi, inferne foliosi, in pedunculum longissimum, simplicem aut ramosum, soluti. *Folia* impari-pinnata, foliolis ovali-lanceolatis, mucronatis, integerrimis, junioribus subvis praecipue pubescentibus. *Stipulae* subulatae, apice subsetaceae.

Habit. in Chilensi Regno. 21.

399. *HIPPOCREPIS annua*: pedunculis umbellariis subtrifloris: leguminibus circularibus altero margine sinuatis ad semina villososacris: foliis pinnatis lineari-oblongis.

Venit locis montosis aridis et inter saxa en *Velaz Rubio* oppido Regni Granatensis, circa Orcehim: Murcia urbem, alibi in Hispania meridionali.

Floret ab Aprilis mense ad Augustum. 22.

300. *ORNITHOPUS nudiflorus*: foliis impari-pinnatis petiolatis: pedunculis nudis subtrifloris: leguminibus cylindricis incurvis: caule diffuso procumbente. *Legas. Variet. ann.* 2: num. 19. p. 40. *Pers. loc. cit.*

301. *Pusillum ornithopodium* Pena. *Dalech. Hist.* 487.

302. *Ornithop. ebracteatus*. *Bror. Fl. Lusit.* 2. p. 159.

Legi in Principatu Asturicensi ad cras fluvii *Nabalon*, prope Pravia oppidum: cel. *Cavanilles* in Regno Valentino in tractu *Vall de Jenes*.

Floret Aestate. 23.

303. *DALIA virgata*: decandra, foliis multijugis: foliolis ellipticis retusis: spicis ovatis terminalibus subaggregatis: dentibus calycinis hirtis.

Habitat in Imperio Mexicensi.

In H. R. M. Aestate et Autumno floret. 24.

304. *DALIA tuberculata*: spicis cylindricis: ramis virgatis tuberculatis: foliis canescentibus pinnatis, foliolis obovato-cuneatis emarginatis.

*Parosella tuberculata*. *Blench. H. R. M. ann.* 1805 et 1816.

*Dalea mutabilis* Pers. admodum affinis, sed notis enumeratis, spicisque gracilioribus satis diversa.

Habit. in Nov. Hisp. Semina adportavit prael D. *Sessé*. 2.

305. *DALIA sericea*: spicis oblongis densis sericels longissime pedunculatis: foliis impari-pinnatis, foliolis omnibus ovali-lanceolatis, subvis praecipue sericels.

Hab. in N. Hisp. 25? Pili sericels ferruginei. *Spica Trifolii* angustifolii similima.

306. *TRIPOLIUM clandestinum*: capitulis ovatis terminalibus axillaribusque pedunculatis: calycibus rigidis inaequalibus: corollis clandestinis: foliolis obovato-cuneatis pubescentibus subtintegerrimis.

Venit locis rudertis circa Matritum.

Floret Mayo et Junio. 26.

307. *TRIPOLIUM conicum*: capitulis axillaribus pedunculatis oblongo-conicis densis: calycibus subaequalibus dentibus subulato-setaceis: foliolis obovato-cuneatis emarginatis integerrimis pubescentibus.

Habitat

308. *TRIFOLIUM serrulatum*: capitulis umbellariis globosis, fructiferis reflexis: calicis dentibus inaequalibus corollae subaequalibus: leguminibus dispersis: foliolis oblongo-obovatis serrulatis. — Juxta Trifol. angulatum: *Wulderst.*

H. locis arenosis circa Matritum.

309. *LOTUS rubriflorus*: hirsutus: caule ramoso. diffuso: pedunculis 2-3-floris: leguminibus erectis teretibus corollisque calyce longioribus. *Lag. Variet. num.* 22. pag. 213.

*Pers. Encir. bot.* 2. p. 354.

310. *Lotus microcarpos* Brot. Fl. lus. vol. 1. pag. 2.

Habitat prope Praviam oppidum, in Asturum Principatu, locis arenosis.

Floret Julii et Augusti mensibus. 27.

311. *LOTUS rubripinnatus*: hirsutus: pedunculis axillaribus unifloris brevissimis: foliolis lanceolatis: stipulis foliaceis solitariis: leguminibus rectis. — Pone *Lotum tetraphyllum*. Lian.

Habit. in *Talesbanano* in Chilensi Regno. 28? *Foliola* in petioli communis apice gemina, tertium sub apice, vel ad petioli medium. *Stipulae* foliaceae, solitariae, ovato-lanceolatae. *Legumina* hirta, calyce longiora.

312. *LOTUS trichocarpus*: pedunculis unifloris petiolo longioribus: leguminibus falcatis longissimis tomentosis. — Pone *Lotum contimbriense*. Pers.

*Planta* herbacea, decumbens, subpedalis. *Foliola* cuneata, pilis raris albis molibus, subvis praecipue, marginibusque obsita, ut bractee et stipulae. *Stipulae* cordatae obliquoae, petiolo communis duplo longiores. *Pedunculi* axillares, pilosi, apice geiculati, inflexi, ibidemque. *Bractea* subsessili, triphylla suffultis: foliolis ovatis, calycis longitudine. *Corolla* ex albo-pallens, calyce longior. *Carina* acuta, apice atropurpurea. *Legumen* subtriloculare, teretiusculum, torulosum, tomentosum, falcatum.

Patriam ignoramus. In Herbario *Clar. Camerilles* specimen unicum hujus plantae, cum aliis *Loti contimbriensis*. Brot. in Valentiae Regno lectis aservatur: sed cum planta enata



sit in eadem olla, in qua anno proximo elapsa Daleae species sata fuerit, de patria dubius haereo.

310. *GLYCYME lanceifolia*: caule volubili scandente: foliis ternatis: foliolis angustolanceolatis glabris, lateralibus obliquis: pedunculis axillaribus solitariis racemoso-quadrifloris.

Habitat in Canariis?

In H. R. M. floret Vere. 7.

## CLASSIS XVII. SYNGENESIA.

### ORD. I. POLYG. AEQUALIS.

#### \* *Cicboraceae. Juss.*

311. *PICTIDUM pinnatifidum*: foliis amplexicaulibus oblongis pinnatifidis albo punctatis: pedunculis squamosis incrassatis: anthodis squarrosis.

*Sonchus hispanicus. Jacq. Hor. Schoem. 2. pag. 9. tab. 143.*

Venit solo arenoso inter sata circa Malacam et in Valentiae Regno.

Floret Vere. 6.

312. *SONCHUS simplicissimus*: pedunculis axillaribus squamosis paucifloris folio brevioribus: foliis oblongis dentato-spinosis: caule simplicissimo.

H. prope *Quintanar de la Orden, Aragon* oppida, alibique locis argillosis.

Floret Julio. 2.

313. *SONCHUS scorzoneraeformis*: pedunculis squamosis unifloris: squamis anthodii margine membraceis: foliis pinnatis, pinnis linearibus decurrentibus.

Venit in collibus aridis cretaceis circa Heliocratam, Murciam urbem et alibi in Murciae et Valentiae Regnis. — Floret Vere. 2.

314. *PEDANTHERUS ruscifolia*: foliis radicalibus ramulosis spinuloso-ciliatis: ramis divaricatis bifidis: floribus geminis solitariisve suboppositis pedicello squamoso longioribus. — *Sonchus reflexus. El. p. 15.*

Habit. ad saxorum calcareorum rimas in Monte Sancti Michaelis juxta Orcehim, Alonnam, *Cuevas Overa* oppidum, alibique in Hisp. meridionali. Floret primo Vere. 2.

315. *THRINCIA laevis*: radice fasciculata: foliis, scapis, pedunculis, anthodisque laevibus glaberrimis.

Habit. in Hispania.

316. *THRINCIA nudicalyx*: calycibus glabris laevibus subdodecaphyllis: foliis lanceolato-oblongis sinuato-dentatis hirtis, pilis furcatis.

*Radix* simplex: folia obtusa, nonnunquam fere runcinata, subtus praesertim pilis furcatis hirtula.

Venit locis lapideis, saxosisque circa Orcehim.

Floret Januarii et Februarii mensibus. 6?

#### SOLDEVILLA. Lag.

*Char. gen. different.*

*Anthod.* monophyllum, fructiferum basi ventricosum, lacinialis tunc in coam convergentibus. *Recept.* paleaceum, paleis multido-setaceis: *Pappus* nullus.

317. *SOLDEVILLA setosa. Lag. Varied. de*

*Climax. vol. 2. in 19. Pers. Enchyr. bot. 2. p. 348* ubi *calyx*: communis perperam dicitur *imbri-catus*. — *Hispidella hispanica. Barnades secundum. Cav. Herb. — Lam. Encycl. 3. p. 134.*

Hab. in locis arenosis aridis in utraque Castella. 6.

#### \*\* *Cynarocephales Juss.*

318. *CARTHAMUS hircinus*: caule erecto hirsuto subbifloro: foliis amplexicaulibus pubescentibus ovato-lanceolatis sinuatis spinoso-dentatis pungentibus.

*Carth. arborescens. Linn?*

Venit ad montium radices circa Malacam, Orcehim, Murciam urbem, alibique in Murciae, Valentiae et Granatae Regnis. — Odorem hircinum spirat.

Floret ultimo Aprilii et Majo. 5.

Vulgo *Gordo cabra* Orcehitanis audit.

319. *CIRSIUM cernuum*: foliis subamplexicaulibus lanceolatis sinuato-pinnatifidis spinosis (subtus arachnoides): calycibus arachnoides-tomentosis cernuis.

*Cirsium cernuum. Elench. H. R. M. ann. 1805.*

Habitat in Imperio Mexicano. 2.

320. *CIRSIUM horridum*: arachnoidem: caule erecto subtrifloro: foliis semiamplexicaulibus spinosis bifariam pinnatifidis, lacinis profundissimis lanceolato-subulatis, integris: calycibus longe valdeque spinosis. *Lag. Varied. ann. 2. n. 22 pag. 214.*

Legi in Granatensi Regno juxta viam, quae ab *Albama* ad *Velez-Málaga* ducit, prope la *Venta alta*. — Floret Autumno. 2. — *Corgia alba.*

321. *STRAELINA pinnata*: tomentosa: foliis pinnatis, pinnulis linearibus decurrentibus margine revolutis: pedunculis unifloris.

Habitat in Extremadura, *Maacha* et Granatensi Regno, alibique in Hispania meridionali.

Floret Junii, Julii et Augusti mensibus. 2.

#### \*\*\* *Discoidaeae. Linn.*

#### G. N. PODANTHUS.

*Ord. nat. Corymbiferae Juss.*

*Char. gen.*

*Anthod.* subimbricatum, parvum, multiflorum. *Flosculi* pedicellati, 5-dentati, aequales. *Recept.* paleaceum. *Pappus* coronula dentata.

322. *POD. ovatifoliae. Frutex* ramis subangulatis, trichotomis, junioribus pubescentibus. *Folia* opposita, breviter petiolata, ovata, acutiuscula, scaberrima, subtus subpubescentia. *Pedunculi* terminales, terni, uniflori, subpollicares, dense villosi, in fructu elongati sesqui-bipollicares.

Hab. circa *Concepcion* urbem, et prope *Chilian* oppidum Regni Chilensis, ubi vernacule *Palo negro* nuncupatur, uti ex scheda cel. *Ludovici Néé* constat.

#### G. N. CALYDERMOS.

*Ord. nat. Fus. Corymbiferae.*

*Char. gen. different.*

*Anthod.* oblongum, imbricatum, scarlosum. *Corolla* floeculosa, flosculis 5-dentatis.

*Recept.* paleaceum. *Semina* uniformia, turbinata, caute.

333. *CALYD. scaber*: foliis ovato-lanceolatis utrinque scaberrimis.

Planta herbacea, subpedalis, perennis, scaberrima. *Caule* teres, erectus, hispidulus, plerumque simplicissimus. *Folia* opposita, sessilia, vel subsessilia, subcordata, ovato-lanceolata, vel ovata, acuta, remote dentato-serrata, tripinnaria, utrinque, sed sub-ter praecipue villosa scabro obducta. *Umbellae* terminalis, longe pedunculata; *radix* tres tomentosa, floribus fasciculato-cymosis. *Pedunculus* folio uno, duobusve alternis, stipitatus. *Pedicelli* breves, bractea parva suffulti. *Calycis* squamae lutescentes.

Hab. in *Santa Monica*, *Guanaquato*, *Acapulco* et *Real del Monte* oppidis Imperii Mexicani. L. (V. S. Fl. fr.)

334. *CALYD. longifolius*: foliis oblongo-lanceolatis laeviusculis.

a foliis oppositis.

a foliis verticillato-ternis.

*Caule* erectus, teretiusculus, subpilosus, nitidulus. *Folia* oblongo-lanceolata, acutissima, sessilia, remotissime dentato-serrata, subpilosa, laeviuscula, supra nitida, subtripinnaria, ad basin 3-4-lineae lata. *Pedunculus* terminalis, longissimus, subpedalis, apice umbellatus, radice tribus quatuorve, compactiusculis, apice fasciculato-corymbosis. *Inflorescentia* 1-3-phyllum. *Foliolis* linearibus, integerrimis. *Pedicelli* foliis brevioribus suffulti.

Hab. in N. Hsp. 2.

#### G. N. THYMOPHYLLA.

*Ord. nat. Corymbiferae. Juss.*

*Char. gen. diffinit.*

*Anthod.* monophyllum, campanulatum, dentatum, multiflorum. *Recept.* nudum. *Pappus* pentaphyllus, paleaceus, paleis truncatis, brevibus.

335. *Thym. retifolia*. *Rami* subfiliformes, articulati, oppositi, foliis numerosis, confertis vestiti. *Folia* opposita, sessilia, vix sexquilineam longa, tomentosa, subsetacea. *Pedunculi* terminales, elongati, villosuli, folio uno alterove ornati, rigiduli; bipillicares uniflori. *Calyc.* extra pulverulento-tomentosus. *Corolla* atropurpurea, ut in sicco apparet. *Semina* longitudine calycis.

Hab. in N. H. Suffrutes.

#### G. N. MICROSPERMON.

*Ord. nat. Corymbiferae. Juss.*

*Char. gen.*

*Anthod.* campanulatum, sequele, multiflorum. *Pedicelli* radii 6-8 rariore, limbo sublobulato. *Recept.* nudum. *Pappus*, *Denticuli* brevissimi et aristae 1-3.

336. *Microsp. subulicinctum*. — *L.* *Levis* decumbens: caule simplici decumbente, foliis subcordatis, obtusis, serratis. *Fructus* fl. ind. occid. 3. pag. 1221? — *Flora. Jacq.* 180. pag. 288. *tab. 122. fig. 2.* *Spina* campanulatum nostrum ut bene exprimit. *Pedicelli* lobulati, vix triphylli. *Calyc.* filiformis, decumbens, lobulatus, simplex. *Inflorescentia* 1-3-phyllum. *Foliolis* linearibus, integerrimis, bipillicares uniflori. *Calyc.* extra pulverulento-tomentosus. *Corolla* atropurpurea, ut in sicco apparet. *Semina* longitudine calycis.

brevissime petiolata, crenata, rotundato-subcordata, frequentius rotundato-ovata. *Pedicelli*. Flore multo longiores.

Habit. in *Guanaquato*, *Queretaro* et *Salvatierra* in Nova Hispania. L.

337. *SANTOLINA viscosa*: suffruticosa: ramis viscidis: foliis inciso-pinnatis filiformibus: floribus subcorymbosis.

Habit. in collibus aridis sterilibus circa *Cuevas Overa* oppidum, ad saxorum rimas juxta *Albama*, alibi in Murciae Regno.

Floret Aestate et Autumno. 2.

338. *SANTOLINA pectinata*: pubescens: foliis junioribus teretiusculis tuberculatis postea pinnatifido-pectinatis, ramis unifloris.

Leucanthemum fruticosum pinnatis superioribus Quamoclit foliis, limbo verniculatis et torosis. *Barrst. obs.* 1103. tab. 428. bona: plantam innotam representans.

Habit. in Regno Granatensi, circa *Albes* et *Puebla de D. Fadrique* oppida legit col. D. Clement.

339. *SANTOLINA canescens*: tomentoso-lucans: foliis inferioribus quadrifariam tuberculato-dentatis, pedunculorum linearibus planis integris.

Abrotanum flemius IV. *Char. Hist.* 1. pag. 341.

*Pedunculorum* folia aetate fere glabra, obtusa, nec acuta ut in *Santol. cormaripifolia*. Linn.

Hab. in Hispania meridionali. Collitur subdio in R. M. H. ubi Junio et Julio floret. 2.

340. *CHRYSOCEMA verticillata*: anthodis squarrosis: foliis lanceolatis sessilibus verticillis.

Venit ad rupium fissuras, circa *Alouam*. *Orcellini*, *Murcia* urbem, alibi in Murciae, Graustae et Valentiae Regnia.

Floret Autumno. 2.

341. *SPARGANOPHORUS ageroides*: corymbis terminalibus: foliis ovatis acutis serratis tripinnaribus.

*Anthodii* squamae, erectae, lineares, acutae. *Corolla* caerulea.

Hab. in N. Hispania. 2. *Semina* adportavit, *praesent.* D. *Jard.* ann. 1804.

342. *SPARGANOPHORUS obscurifolius*: foliis oppositis petiolatis oblongo-lanceolatis obtusis crenatis.

*Corymbis* terminalis, compactus, longissime pedunculatus. *Corolla* ex albo-caerulea.

Hab. in N. Hsp. 2.

343. *EUPATORIUM micranthum*: foliis ovato-lanceolatis acutissimis serratis, basi cuneata apiceque integerrimis.

Juxta *Eupatorium caelestinum*. L.


Planta fruticosa, scaberrima, subtripinnaria. *Corymbi* fasciculati, terminales. *Anthodii* subimbricati, multiflori. *Corolla* caerulea.

Hab. in Nova Hispania. 2.

344. *EUPATORIUM secundiflorum*: fruticosum: foliis cordato-oblongis crenato serratis scabris trinerviatis: floribus cernuis in racemum terminalem foliorum digestis.

*Anthod.* oblongum, cylindricum, apice imbricatum, squamis lanceolatis subulatis, apice late mucronatis. *Corollae* lobulatae lobes obtuse quinque, albo, vix calycem superante, patulo, in anthera connatis. *Stigm.* lanceolata, elongati, originato-crenatis.





cula, obtusa, apice crasiore rufa-fusco. Anthodis intra tum.

Hab. in N. Hispania. — Floret à Februario ad Aprilim mensem in intern. M. R. M. II.

# PALAFOXIA.

Ord. nat. Corymbiferae. Just.

Char. gen. contractae.

Anthodium oblongum, subimbricatum, 8-polyphyllum, multiflorum, in fructu stellato patens. Corolla flosculosa, calyce longior; flosculis quinquefidis. Pappus polyphyllus, paleaceus. Receptaculum nudum. Semina marginalia foliolis calyculi patentibus involuta.

335. PALAFOXIA linearis. (L.) Presl. n. 464. — Ic. vol. 3. p. 3. tab. 305. Agerati nomine.

Palafoxia genus Steviae et Agerati vicinum, sed diversum ab Stevia, calyce oblongo subimbricato, multifloro, in fructu stellato patente, foliolis semina amplexantibus: atque pappi paleis lanceolatis, acutis. Ab Agerato calyce oblongo, corolla duplo breviori, in fructu stellato patente, atque flosculis limbo quinquefido distinguitur.

# STEVIA. Cavan.

Char. gen. different.

Anthodium tubulosum, simplex, pentaphyllum, quinqueflorum, corolla duplo vel amplius brevis, immutatum persistens. Corolla flosculosa, limbo 5-fido. Receptaculum nudum. Pappus paleaceus, paleis brevibus, muticis, vel longe aristatis.

Char. Pappi paleae breves, truncatae, 5-8, in paucis muticae omnes, frequentius vero earum nonnullae, rarissime omnes apice inagristem calyce longiore, crassissime denticulata desinentes.

Obs. 1. Stevia connata anthodia nonnulla gerit 6-phylla. 2. 7-florea.

# Herbaceae.

336. STEVIA serratae foliis linearilanceolatis, glabrisculis serratis, basi integerrimis: in petiolum attenuatis, subsecundis, pappo triaristato.

a. Stev. serrata: caule glabrescens, foliis glabris. Cav. Ic. vol. 3. pag. 33. tab. 355.

a. caule subhirsuto, foliis subtus pilosis latioribus sublinearilanceolatis.

Variet floribus albis et ex albis purpureiscentibus.

Hab. in N. H. II.

337. STEVIA glandulosa: pappo paleaceo triaristato: foliis linearibus subsphaelatis apice dentatis, in petiolum attenuatis superioribus linearibus subintegerrimis: caule pubescente simplici rectifloro.

Stev. pubescens. Elench. H. R. M. ann. 1805.

Stevia purpurea. Hornem. H. R. M. p. 302.

H. in N. H. — Circa Guanauro legitur Cel. D. L. N. II.

Causteres, pubescens, subhirsutus, erectusculum, simplex superne tantum ramulis nonnullis corymbiferis. — Folia inferiora opposita, lanceolata apiculata, in petiolum angustata, superne dentata, reliqua sparsa, subintegerrima, linearia, versus apicem paulatim latiora. Corymbus flagitia tunc densiusculi.

Anthodia pubescentia. Corolla purpureo-sanguinea. Pappus paleis tribus truncatis brevibus, cum quibus totidem aristae alternant.

Hab. in N. H. — Circa Guanauro et Chama oppida legitur Cel. Peregrinator L. N. II. — Semina communicavit praed. D. Stev. Agerati nomine.

338. STEVIA lanceolata: foliis lanceolatis acutis planiusculis in petiolum attenuatis serratis subtus scabris, summis linearilanceolatis integerrimis: pappo paleaceo triaristatoque.

Ageratum Sessé.

a. floribus albis, foliis subtus glabris.

b. floribus purpureis, foliis subtus subhirsutis.

Caulis 2-3 pedalis, subhirsutus, corymborum poudere tandem decumbens, superne ramoso-corymbosus. — Folia lanceolata, utriusque acuta, primo cymbiformia, deinde planiuscula, tripplinervia, basi integerrima, in petiolum attenuata, exinde serrata, serraturis acutiusculis, rames et floralia linearilanceolata, integerrima, aut subintegerrima. Corymbus flagitia tunc laxiusculi, multiflori. — Villositate longiori, statura longe majori, foliisque lanceolatis serratis triplo fere latioribus quam in Stev. pubescente ab hac et hyssopifolia abunde differt.

Hab. in N. H. Circa Pachma oppidum legitur Cel. Lud. N. II. — Semina adportavit D. Sessé ann. 1804.

339. STEVIA purpurea (quinquearistata): pappo paleaceo aristatoque, aristis quinque.

Stevia purpurea. Elench. H. R. M. anno 1805. — Ageratum Sessé.

Diversa ab Stev. purpurea. Pers.

Planta herbacea, pubescens. Folia caulicea sublinearia, cymbiformia, lasima, opposita, superne dentata, reliqua sparsa, angustiora acuta, subintegerrima. Flores laxi corymbosi, pedicellis unibifloris. Corolla purpureo-sanguinea. Pappus constanter quinquearistatus.

Hab. in Irmiguigan, Sta. Monica et Guadalupe in N. H. ubi legitur D. L. N. II. — Semina communicavit D. Sessé ann. 1804. Agerati specie nomine: Monch. 29. ann. 1805.

340. STEVIA pilosa: pappo paleaceo aristatoque: caule diffuso subpubescente: foliis linearilanceolatis superne dentato serratis supra lucidis sessilibus.

Stev. hyssopifolia. Cel. D. L. N. II. ann. 1805.

Ageratum punctatum. Ort. Decad. p. 17.

Similis Stev. pubescenti, sed caule diffuso brevissime pubescente, foliisque sessilibus confertioribus, superne lucidis, subhirsutis, atque corymbo confertiori satis diversa. Pappus 3-5-aristatus.

Obs. Nedium haec, sed et reliqua omnes Steviae species folia utrinque punctata praesunt.

Hab. in Regn. Mex. II. ann. 1805.

341. STEVIA pilosa: pappo paleaceo aristatoque fasciculato corymboso: foliis sublinearibus dentatis, caule piloso.

Stevia pilosa. Elench. H. R. M. ann. 1805.

Ageratum Sessé.

Caulis teres, subhirsutus pilis viride albis virentibus, superne ramosus. Rami conimiles erecti corymboso-flagitia, terminati corymbis fasciculato corymboso multifloris. Folia

lia sublinearis, cymbiformis, superne dentata, sparsa, infima opposita. Calyces purpureo-centes. Flores sanguinei. Pappus paleaceus, muticus.

Hab. in N. H. Circa Guanajuato legit prael. D. L. Née. — Semina communicavit D. Sesse.

342. STEVIA *subulosa*: pubescens foliis inferioribus ovatis crenato-serratis; superioribus sublinearibus serratis, summis subintegerrimis; pappo 3-aristato.

Planta tota pubescens, odorem gratum spirans, subtripedalis, sequenti sat similis. Folia caulina et ramea inferiora opposita, petiolata, ovata, crenato-serrata, basi cuneiformi integerrima, triplinervia, ramea superiora alterna, ex basi cuneiformi ovato-lanceolata, lanceolataque, floralia angustiora sessilia integerrima, vel denticulo uno alterove notata. Corolla sordide alba, raro laete rosea. Pappus paleaceus, 3-aristatusque, nonnumquam seminibus duobus muticis.

Hab. in N. H. Circa Guanajuato legit D. L. Née. Semina communicavit prael. D. Sesse.

343. STEVIA *paniculata*: caule paniculato: foliis inferioribus ovatis, superioribus ovato-oblongis serratis basi cuneatis integerrimis, summis linearilanceolatis; pappo mutico subaristatoque.

Stevia paniculata. Elench. H. R. M. ann. 1805.

Caulis teres, erectus, superne ramoso-paniculatus, breviter pubescens. Folia caulina inferiora opposita, petiolata, ovata, acuta, serrata, basi cuneata, integerrima, triplinervia; superiora sparsa, inferioribus similia, sed basi longius angustata, breviusque petiolata; superiora sessilibus angustiora, minus dentata, sessilia, floralia integerrima linearilanceolata. Pedunculi filiformes, villosi, stricti, apice bifidi, 3-4 flori, corymbum constituentes. Corolla limbo albo, tubo ex albo sordide purpureo-centes. Semina tria pappo paleaceo mutico, duo reliqua pappo paleaceo 1-2-3-aristato.

Hab. in N. H. ubi legit D. Lud. Née. Semina adportavit D. Sesse ann. 1804.

344. STEVIA *ovata*: foliis ovatis triplinerviis serratis, basi cuneatis integerrimis; corymbis fastigiatis compactis; pappo paleaceo subuniaristato.

Stevia ovata. Elench. H. R. M. ann. 1805 — *Hornem. loc. cit. p. 793.*

Planta herbacea, valde ramosa, glabrescens, subquadripedalis. Folia pleraque opposita, deltoides-ovata, vel si mavis, ovata basi cuneata. Corolla purpurea limbo dilutiore. Pappus paleaceus, in seminibus nonnullis uniaristatus.

Hab. cum praecedentibus. Semina missa praelud. D. Sesse. 21.

345. STEVIA *rubroaristata*: pappo 7-9-aristato.

Habitu Steviae ovatae similis.

Hab. in Peruviae Regno. 21. D. Ludovicus Née legit.

346. STEVIA *enarthotricha*: panicula trichotoma ramosa, pedicellis unifloris; foliis caulinis oppositis ovatis, utrinque acutis serratis, petiolatis; pappo subtriaristato.

Caulis hirsutus, hirsutie articulata, tandem omnino fere deponenda, superne in paniculam amplam solutus. Folia 3-plinervia,

caulina opposita, petiolata, ovata, utrinque acuta, basi cuneata, integerrima, ex inde dentibus latis obtusis serrata, supra praesertim pilis raris subhirsuta, superiora fere lanceolata; ramea inferiora opposita, subintegerrima, reliqua pedunculorumque sparsa, integerrima; parva subuliplica. Pedicelli uniflori, anthodio glabro acuto longiores. Corolla purpurea. Pappus paleaceus, uniaristatusque.

Hab. in N. H. 21.

347. STEVIA *adenophora*: panicula trichotoma laxa ramosa; pedicellis unifloris villosis glanduliferis; foliis caulinis oppositis sessilibus ovatis obtusis serratis.

Planta herbacea, perennis, villo brevi glandulifero scatenis. Folia lato-ovata, obtusa, triplinervia, incisio-serrata, serraturis obtusis, basi subdeltoides integerrima, sessilia; ramea inferiora opposita, multoties minora serrata, floralia seorsim minora integerrima. Panicula ut in *St. enarthotricha*. Calyx villosus. Corolla purpurea. Pappus triaristatus, raro aristis quatuor, vel paucioribus.

Hab. in Regno Chilensi, ubi legit D. Ludovicus Née. 21.

348. STEVIA *incanescens*: foliis crenato-serratis subtus tomentoso-incanis.

Stevia incanescens. Elench. H. R. M. ann. 1805.

a. pappo paleaceo-aristatoque, aristis quatuor.

b. pappo paleaceo uniaristatoque.

c. pappo paleaceo, mutico.

Hab. in N. H. 21. Semina adportavit D. Sesse.

349. STEVIA *micrantha*: foliis ovatis subcordatis triplinerviis serratis oppositis alternisque; pedunculis subunifloris; pappo triaristato corolla longiori. — Stev. micrantha. Elench. H. R. M. ann. 1805.

Planta annua, vix pedalis *Agerati conizidae* Lim. facie. — Caulis, rami, petioli, pedunculi et calyces villosi. Flosculi 5, albi, calyce duplo longiores; limbo minuto 5-dentato, rarius 4-dentato. Pappus aristae tres, setaceae, flosculis paulo longiores, creberrime denticulatae, basi dilatatae, cum quibus totidem paleae brevissimae, truncatae, alternant.

Hab. in N. H. 21. Semina communicavit cl. Sesse. *Agerati* nomine. D. Ludovicus Née legit circa Guanajuato.

350. STEVIA *trifida*: foliis triindis linearibus.

Herbacea. Caulis filiformis debilis, purpureascentis, pube brevissima, glandulifera. — Folia linearia, angusta, caulina opposita, trifida, laciniae linearibus, intermedia longiori ad medium dente utrinque plerumque excisa, floralia alterna, indivisa. Corymbus pauciflorus terminalis. Anthodium villosissimum, foliolis acuminatis. Corolla purpurea. Pappus paleaceus triaristatusque.

Hab. in N. H. (V.S.) 21?

\*\* Fruticosa.

351. STEVIA *connata*: foliis connatis oblongo-lanceolatis acutissimis serratis; caule stricto; pappo paleaceo aristatoque.

Stevia connata. Elench. H. R. M. ann. 1805. Planta fruticosa, glaberrima, ergyalis, et ultra. Folia superiora nonnumquam sparsa.

*Gymnema terminalis*, Burtholm, dentata, foliis alba. Ceylon p. 6-7. Hort. — *Peltandra* canaliculata et canaliculata stria distincta. H. in N. H. 2. (VV.) *Stemina communis* vix D. dard.

352. *STEVIA subpubescens*: pappe palencae mucico: foliis ovatis-lanceolatis subtus pubescentibus.

*Caule* fruticoso. Rami teretes, superius pubescentes, uti pedunculi, pedicelli, et florum pagina inferior. Pedicellipappi, pagina cernua longiora, ovata-lanceolata, apice obtusa, obscure remotiusque dentata. Corymbi cernuique fastigiatos et corymbis cernuique multiflori. *Anthed.* subvillosus. *Cavendish* alba-carnet. Pappus paleaceo mucosus.

Hab. in N. H. 2.

353. *STEVIA salicifolia*: foliis lanceolatis apicatis, ad medium remote serratis: pagina cernua.

*Stem salicifolia*. Cav. Don. vol. 4. pag. 32. tab. 334.

*Aperum* viscosum. Cav. Don. p. 34. *Folia* brevissima petiolata, angusta-lanceolata, stylique, apicatis, acuta, integerrima, vel dentibus distis suberrata, opposita cernua, floralibus exceptis. *Petiolus* cernui. Pedunculi villulosi. Pappus paleaceo, argenteo, nonnunc mucosus.

Hab. in N. H. 2. *Stem salicifolia* p. 2.

354. *STEVIA laeta*: pappe palencae mucico: foliis oppositis lato-lanceolatis dentatis lucida.

*Planta* fruticosa, glandulis succulentibus humorem viscidum, quo veluti nititur et lucida, undique obecta. Similis *St. subpubescens*, sed magis humoreosus et abunde diversa.

Hab. in *Impatiens* et *Chimaphila* in N. H. ubi legit D. A. No. (V6.) 2.

355. *STEVIA cernua*: pappe palencae mucico: foliis lato-lanceolatis argute serratis oppositis terneve subcarnata.

*Pedunculi* communiterminati, apice umbellati, radiis 3-5 apice corymbosa, corymbis multifloris, cernuique, foliis. *Folia* ovata ad pedunculorum pedicellorumque pagina cernua, acutissima integerrima.

Hab. in *Portulaca* Regni, ubi legit D. Ludovici No. 2.

356. *HYMENOPAPPUS pedatus*: foliis petiolatis digitatis-pedatis, summis indivisis anthed decidua. — *Hymenop. pedatus*. Cav. Don. — *Stem pedata*. Cav. L. vol. 4. p. 32. tab. 356. *Villosa* p. 2. p. 1775.

Hab. in *Stem* p. 2. p. 793. — *Aperum* pedatum. Cav. Don. p. 32.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

Hab. in N. H. 2. p. 3.

**SECRET**

**JUNE 22, 1954**

1. The first of these is the fact that the  
2. Government has been unable to secure the  
3. necessary funds to carry out its policy.  
4. The second is the fact that the Government  
5. has been unable to secure the necessary  
6. funds to carry out its policy.

1913. *Fagus subtilis*: *folia pinnatis*.  
*stipitata*: *folia lanceolata serrata*: *hor-*  
*tus. lunc corymbosa*. — *Eichh. M. R. M.*  
*vol. 120.*

2. Habitat in *Impatiens mexicana*. C. Florula  
indistinctly D. H. H. an. etc., on combination  
D. *ferax* siliqua.

**Business & Gov. Press.**

WILSON, C. J.

SECRET

**RECEIVED**

**Bureau of Field Inspection**

**Adaptation:** New, Enchirid.

**Page(s) Specimen / amount**

*Pyrenaria* sp. Com.—Willd.

Ord. nat. *Corymbiferae*. *Yucca*. Junta Fago-

**Chemical Abstracts**

...antha, duplex, utrinque octophyllum, interia tubulosa, stamina longius. Recept. glaberrima, densissima. Pappus paleaceus; pappi in tres glaucae incisus, subulosis, densissimis.

34. — *Fraxinus diversifolia* var. *foliolosa* (Mill.) B. S. P. — *Dryas* *glandulosa*. Chen. *Fraxinus* de la plant. p. 309. *Fraxinus diversifolia* Willd. *sp. pl.* vol. 3. p. 489. — *Fraxinus parvifolia* Fenzl. *plant. nov. acc.* at fol. 26.

Occurrit per totumque Sub Americam:  
vulgatissimum est in Sanctaefidelis Regno, ubi  
venerabile fides, in est fides, et gravem adeo  
venerabile fides, et in fidei puerum. (2)

1. 24. Dors. pubescens; caule patula pubes-  
centia; foliis glaucis linearibus apice latioribus  
lanceolatis; pedunculis unifloris; an-  
theris brevioribus.

*Aster pinnatus*, Cav. Ic. 3. p. 6. tab. 212.  
Hb. in M. M.

*Aspidium adnigrum*, *minime imbricatum*.  
*Aspidium* et in *Dact. chrysanthemoides*.

364. *Dracopis subnigerrimus*: foliis pinnatis, foliolis pinnatis acutis canaliculatis subnigerrimis; involucris uniloricis.

-... *Planta, pilosa, subhirsuta. Folia longiora quam in praecedenti, duploque fere angustiora, acuta, serrata, pinnatimque integerrima, rare saepe densa uno dentibus cincta.*

367. *Dryasida togetheri*: foliis cauliculis oppositis, pinnatis linearibus, superioribus serratis; pedunculis umbellatis, calyce exteriori linearis-sulcato.

Planta herbacea Tugalis semibulosa, Caesp. - Caulis teres, striatus, subpubescentis. Folia opposita subulna, lineata, planiuscula, ciliolata, ad mediumnimum superiorem densibus ciliis ciliata, acutis serrata; ad inferiorem repanda, ciliata longioribus; infimis brevibus, demum ciliatis, longioribus, apice setula appendiculata; ciliis ciliata - ciliis - ciliata.

**THE** **NEW** **YORK** **PUBLIC** **LIBRARY** **ASTOR** **TENNYSON** **TRUST**

Hab. in N. Mex.

368. *Drac. appendicularis*: foliis impari-pinnatis; folioli quinque-ovato-cuneolatis inaequaliter arguteque serratis, glandula oblonga apice appendiculatis.

Feb. 1968 Acapulco in W. Mexico, abt  
1000 m. *Peromyscus maniculatus*, D. L. Johnson

Det. Calyce, si cilla exteriori, campanulati, quam frons, corollae, pappe nris cum bracteis, ariscentibus, connatis, omnino: lig. quatuor utriusque media, genere non distinguendos, unius naturaliter vinculo connatis.

356. *Drya. prorepilla*: foliis planiusculis  
dentato crenatis, latisioribus, nervis diaphanis  
hirs. hirsutum densius: ligulis vix calycem su-  
perantibus. *Drya. prorepilla*. Cav. Anal. de  
Cien. nat. vol. 6. p. 334. — Idem. Hort. Borst.  
tab. 62.

4. *Radam rotio orbata*. — *Pteronia pere-*  
*phyllum*. *Cor. Jap. vol.* 2, p. 12, tab. 223. —  
*Widd. sp. plant. Linn. vol.* 1, pag. 1781. —  
*Atroctylis species*, *Yuss. in Anal. du Mus.*  
*de Per.*

Tab. in Nova Hispania. Vidi vivam in R.  
M. Mart. Florit ab Augusto ad Octobrem. ☉

m. *Abies*, *Pinus* et *Quercus* ad Octobrem. C.  
 det. *A. Pterodon* genere, calatim aciculis  
 caryes duplici nec imbricate, corolla radi-  
 ate, et pappo paleaceo-plumbeo: ab *Arctosty-  
 lide*, *Arctostyle* var. *Arctostyle* nat. h. h. h.  
 alis fructificationis characteribus discrimi-  
 nare.

370. *Dios. coccineus*: foliis pinnatifidis dentatis: ligulis obovatis calycem superantibus. — Willdenowia? Gay. Arch.

Tab. propa. *Ismigulipet* oppidum in Nova Hispania. 161

371. *Drym. Cavanillesii*: foliis pinnatis. Sm.

lobis ovatis inaequaliter dentato-sagittatis: paleis pappi interioribus plerumque trifidis.

1. *Adom cista*, *Stenalis* 6-7-8 *Stis*: *trig-*  
matis *trigida*. *Whitewornia glandulosa*. *Cov.*  
*Ann. vol. 1. p. 61. tab. 2.*, *quae fideliter re-*  
*cum a D. Low. in Ill. gen. tab. 66.*  
*Schlechtendalia glandulosa*. *WUHL Sp. pl. vol.*  
*1. pag. 325.* — *Adenophyllum coccineum*.  
*German Revue bot. vol. 2. pag. 458.*

Mat. juxta Dacotinae oppidum in Nova  
Hisp. ubi loci memoratus D. Ludovicus Med.

Obs. Ex unico specimine, quod ex planta  
cultâ in ch. Cavallotti herbario pervenit.

limba et cili. Chlamydomonas spiralis. Limb. 10-12.  
 haec notata digna encerpere placuit. Gamella  
 à petalis marginis et corinis decorantibus sub-  
 angulatus, rufus ramoribus. Petala inferiora  
 opposita, aut subopposita, petalis commutis.  
 Petala opposita, aut subopposita inaequaliter  
 dentata serrata. Pappus 10-14-phyllus, folio-  
 lis aculeatis, immixtus 2-3-lamellis.

1-2 glomeratis, immixtis velidis. 3-4 - ciliatis. 5 - ciliatis. 6 - ciliatis. 7 - ciliatis. 8 - ciliatis. 9 - ciliatis. 10 - ciliatis. 11 - ciliatis. 12 - ciliatis. 13 - ciliatis. 14 - ciliatis. 15 - ciliatis. 16 - ciliatis. 17 - ciliatis. 18 - ciliatis. 19 - ciliatis. 20 - ciliatis. 21 - ciliatis. 22 - ciliatis. 23 - ciliatis. 24 - ciliatis. 25 - ciliatis. 26 - ciliatis. 27 - ciliatis. 28 - ciliatis. 29 - ciliatis. 30 - ciliatis. 31 - ciliatis. 32 - ciliatis. 33 - ciliatis. 34 - ciliatis. 35 - ciliatis. 36 - ciliatis. 37 - ciliatis. 38 - ciliatis. 39 - ciliatis. 40 - ciliatis. 41 - ciliatis. 42 - ciliatis. 43 - ciliatis. 44 - ciliatis. 45 - ciliatis. 46 - ciliatis. 47 - ciliatis. 48 - ciliatis. 49 - ciliatis. 50 - ciliatis. 51 - ciliatis. 52 - ciliatis. 53 - ciliatis. 54 - ciliatis. 55 - ciliatis. 56 - ciliatis. 57 - ciliatis. 58 - ciliatis. 59 - ciliatis. 60 - ciliatis. 61 - ciliatis. 62 - ciliatis. 63 - ciliatis. 64 - ciliatis. 65 - ciliatis. 66 - ciliatis. 67 - ciliatis. 68 - ciliatis. 69 - ciliatis. 70 - ciliatis. 71 - ciliatis. 72 - ciliatis. 73 - ciliatis. 74 - ciliatis. 75 - ciliatis. 76 - ciliatis. 77 - ciliatis. 78 - ciliatis. 79 - ciliatis. 80 - ciliatis. 81 - ciliatis. 82 - ciliatis. 83 - ciliatis. 84 - ciliatis. 85 - ciliatis. 86 - ciliatis. 87 - ciliatis. 88 - ciliatis. 89 - ciliatis. 90 - ciliatis. 91 - ciliatis. 92 - ciliatis. 93 - ciliatis. 94 - ciliatis. 95 - ciliatis. 96 - ciliatis. 97 - ciliatis. 98 - ciliatis. 99 - ciliatis. 100 - ciliatis.





Vault locis ruderalis juxta vias et ad agros  
varietas circa Aloniam, Siciliam, Tarentum  
oppidum, alibi in Murchia et Valen-  
tina Regnis.

Florat Aprilis et Majo mensa. 21.

383. *ACMELLA hirta*: caule foliisque ova-  
lis integris hirtis: radio 12-flo.

Planta herbacea, palmaris, pils albis hir-  
ta, *Cisti guttari*. L. fide. ☉

Hab. in America: cel. Lud. Noë legit.

G. N. MOCHINA.

*Ordine nat. Juss. Corymbiferae juxta Ga-  
lileum Pl. P.*

*Charact. gener. differ.*

*Corollae ovatum, imbricatum. Corolla ra-  
diata, radio paucifloro. Recept. paleaceum.  
Pappus polyphyllus, paleaceus: pappi subim-  
bricatis, lanceolato-subulatis, imbricatis.*

384. *MOCHINA serrata*: foliis oblongo-lan-  
ceolatis serratis.

Planta fruticosa. — Foliis oblongo-lanceo-  
latis, acuta, serratis, mucronata, subtus pubescentia, scabra, breviter petiolatis. Flores  
pedunculati, fasciculati, fasciculis axillaribus,  
terminalibusque.

Hab. in Imp. Mexicano: prope Salaman-  
cam urbem. D. Led. Noë legit.

385. *MOCHINA brachiatæ*: ramis brachiatis:  
foliis ovatis obtuse mucronatis.

Habit. in Peruvia. ? — Flores fascicu-  
li multiflori, glomeris terminalibus.

G. N. VILLANOVA.

*Ord. nat. Corymbiferae. Juss. juxta Va-  
liam.*

*Char. gen. differ.*

*Anthodium hemisphaericum, pentaphyllum.  
Radius 2-10-flores, ligula retundata, brevi-  
bus. Receptaculum minutum, nudum. Semina  
cuneiformia, calva.*

386. *VILLAN. oppositifolia*: foliis opposi-  
tis tripartitis: corollis decem-radiatis.

Hab. in Peruvia. ☉?

387. *VILLAN. alternifolia*: foliis alter-  
nis oppositis: corollis quique-radiatis.

Hab. in Peruvia circa Limam. ☉?

388. *VILLAN. pinnatifida*: foliis ternatis,  
inferioribus oppositis, superioribus alternis:  
foliis pinnatifidis.

Hab. in Portu Roman. ☉? — Cel. Perc-  
grinator in Led. Noë tres hæc *Philenovæ*  
species legit, primam *Ambrosie* nomine,  
posteriores duas innotuitas accepi ex Her-  
bario Cavendishiano.

G. N. FERDINANDA.

*Ord. nat. Corymbiferae. Juss.*

*Dist. inter Anthodium et Anacyclum,  
Charact. genericus.*

*Anthodium hemisphaericum, imbrica-  
tum: lobis subtriplici serie positis, corol-  
la brevibus.*

Corolla composita, radiata. Radius sub  
setulosus. Discus multiflorus

Propter Radii ligulata, ovata, retundatæque,  
pappi tridentati, fascimæ, formis  
pappi infundibuliformis (limbo æquali, quin-  
quedo, cuneato; hermaphrodita, fertili-  
bus).

*STAMINA quinquæ: Filamenta capillaria,  
tubo inserta, libera. Antheræ in tubum con-  
natæ, suberectæ.*

*PISTILLUM: Ovarium utriusque obovatum,  
compressilobum, obscure tetragonum.  
Stylus filiformis. Stigma utriusque bilobum,  
post anthesin erectum, revolutum.*

*SAMEN nudi et disci conformis, obova-  
ta, basi acuta.*

*Pappus in seminibus radii paleaceus, se-  
leis 2-3, brevibus; disci nullus.*

*RECEPTACULUM conicum, paleaceum.  
PREGRAPHUM nullum, propter calyculum  
proprium adhaerentem: Anthodium immu-  
tum, et receptaculi paleae carinatae semina  
fovent.*

*Character generis differentialis.*

*ANTHODIUM hemisphaericum, imbrica-  
tum. — Corolla radiata. Receptaculum palea-  
ceum. — Pappus radii paleaceus, brevis;  
disci nullus.*

*Character habitus.*

*Plantae frutescentes. Folia alterna, opposi-  
tæ, simplicia. Flores corymbati, ter-  
minales.*

FERDINANDO VII, Hispaniarum et India-  
rum Regi Augusto, exoptatissime, Scientia-  
rum naturalium, Botanicae præcipue mun-  
tificentissimo Patrone, hæc novum corymbi-  
ferarum genus, in amori, fidei, et gratitu-  
dinis testimonium, qua maxima possumus  
reverentia consecramus.

*Species.*

389. *FERDINANDA augustæ*: foliis subtus  
argenteis, surculis integerrimis supra præci-  
pue aureo-flavicantibus.

Habitat in Nova Hispania. ?

390. *FERDINANDA emiliae*: foliis subtus  
pubescentibus supra scabris; petiolis folia-  
ceis-alatis.

Habitat in Nova Hispania. (Vidi Siam) ?

G. N. LASTIOFERMUM.

*Locus juxta Anacyclum.*

*Char. gen. different.*

*Anthodium imbricatum, hemisphaericum,  
multiflorum. Corolla radiata, lobulis radii  
obsoletis, integerrimis. Receptaculum palea-  
ceum. Semina globosa, lassa gossypina obo-  
luta. Pappus nullus.*

*Obs. Genus Anacyclo proximum.*

391. *LASTIOFERM. pedunculare*. — *Elench.  
H. R. Mart. ann. 1805.* — Santolinoides al-  
pina saxatilis foliis glaucis et veluti argen-  
teis, floribus luteis. *Michel. Gener. pag. 31.  
tab. 27.* plantam juvenem primo anno flores-  
centem representat.

*Santolina eliocarpa. Pers. Encycl. bot. 2.  
pag. 407.*

*Santolina erecta Linn ? sed synonymon  
Barroetieri plantam in Regno Gratianensi  
crescentem, Santolinæque generi adjuven-  
dam representat, quæ à nostra planta ad-  
modum distat.*

*Planta glauca, glaberrima, laevis subpe-  
dalis, primo anno flores. 21.*

Habitat in monte Morensi in Italia. Co-  
litur subdio in R. Mat. Hort. ubi primo æta-  
tis anno Aestate et Autumno, postea vero ab  
inmate Vere ad Julium mensem flores.

*Obs. Santolina anthemoides. Linn. Flæscules*

radli faemineos fertiles, ligulis obsoletis stipatos promit, ideo ad *Anacyclum* genus amandavimus.

### ORD. 3. POLYGAMIA *frustranea*.

392. *CENTAUREA setifera*: anthodis scariosis; foliis scaberrimis subpinnatis; lacinia linearibus integris margine revolutis, apice setiferis.

Floruit in Reg. Matr. Horto an. 1804 et 1805 ex seminibus à Russia misis. ☉

393. *CENTAUREA sulphurea*: calycibus scariosis ciliatis; squamis exterioribus lanceolatis subulatis; foliis inferioribus pinnatis pinnatifidis subulis arachnoideis, superioribus pinnatifidis lacinia lanceolatis mucronatis integerrimis.

Cent. sulphurea. *Elench. R. M. H. ann. 1805.*

*Corolla sulphurea.*

Ignoratur patria. Colitur subdio in R. M. H. Floret Aestate et Autumno. ☿. — *Centaurea orientalis*. Linn. proxima.

394. *CENT. ligulata*: cano-tomentosa; foliis linearibus ligulatis integris subdecurren-  
tibus; anthodis ciliato-pectinatis.

H. in montibus legionensibus. — Juxta *Cent. montana*. L. locanda. ☿.

395. *CENTAUREA rigida*: anthodis ovatis acutè imbricatis; squamis apice ciliatis; foliis pinnatifidis scabris; lacinia lanceolatis mucronatis, inferiorum lacinia, superiorum pteromque integerrima.

Cent. stricta. *Elench. H. R. M. ann. 1805.*

*Corolla purpurea.*

Ignoratur patria. Colitur in R. M. H. ubi Aestate et Autumno floret. ☿.

396. *CENTAUREA polymorpha*: anthodis ciliato-spinosis, spina terminali patulo-erecta; squamis glabriusculis apice nigro-fuscis; foliis pinnatifidis lyratis subtomentosis. — *Corolla purpurea.*

6. Foliolorum lacinia terminali integra, integerrimaque.

7. Foliolorum lacinia terminali dentata, maxima.

8. Foliolorum lacinia terminali dentata, serrata.

9. Eadem foliolorum lacinia divaricata.

*Centaurea centaureoides* Linn. proxima.  
Hab. in vicis ditiosis *Enenacore* oppidi, alibi in Aragoniae Regno. — Floret aestate et autumno. ☿.

*Fransaladro* vulgo appellatur.

397. *CENT. incana*: tomentosa-lacuna; anthodis ovato-oblongis ciliato-spinosis, spinis recurvatis; foliis pinnatifidis lacinia inferiorum dentatis integris, squamis linearibus.

Venit inter ann. subalpes-catalana, in *Dretille* monte circa Orceolum, alibi in Valentia, Murciae, et Granatae Regna.

Floret Autumno. ☿.

398. *CENTAUREA saxicola*: calycibus ciliatis, squamis interioribus muticis, intermedii spina valida subramosa patente, extimis spinis brevioribus reflexis; foliis pinnatis, summis linearibus.

Habitat ad sanctorum calcareorum rimas juxta Orceolum, alibi in australiore Hispania.

Floret Aprilis et Maja. ☿.

399. *RUSSEZIA incana*: foliis dentatis-ovatis, acutis serratis triplinerviis.  
Hab. in Nova Hispania. Colitur in R. M. H. ubi autumnali et hyberno tempore fleret. ☿. Introducta ann. 1804 ex seminibus per *D. Sord* data.

### ORD. 4. POLYGAMIA *NECESSARIA*.

400. *Microcerus benthicus*: floribus congestis alaribus terminalibusque lina tumida obductis, calycibus subspaspermis.

2. caule erecto simplici aut semel dichotomo. — *Microcerus discolor*. *Pursh. Burser. bot. 2. pag. 425.*

3. multiflorus supinus vel ascendens. *Burser. obs. 99. 12. 1804.*

Habit. in calibus aridis circa Matritium alibi in Hisp. ☿.

401. *Microcerus erectus*: floribus congestis alaribus terminalibusque calycibus linearibus subspaspermis.

Micro. erectus. Linn. — *Leaf. iter. p. 164. tab. 1. f. 3. — Willd. Sp. pl. 2. p. 1385. conclusio Burser. synonyma.*

Hab. cum precedenti. ☿.

### ALGUA. *Cov.*

*Char. gen. differ.*

*Anthodium* simplex, monophyllum, profunde quinque-partitum, patens, mucronatum. *Corolla* radiata, ligulis ovatis, concavis. *Receptaculum* in radio nudum, in disco paleaceum, deciduum. *Semina* corticata. *Pappus* nullus, nisi tubercula seminum pappum dicas.

402. *ALGUA perfoliata*: petiolis basi utriusque appendiculato-filicatis, in pyramidem connatis. — *Alcina perfoliata*. *Cov. B. vol. 1. pag. 10. tab. 15. — Perr. Ench. bot. 2. pag. 491. — Wedelia perfoliata. Willd. Sp. pl. 2. pag. 1335.*

403. *ALGUA ovalifolia*: petiolis connatis inappendiculatis. — *Alc. ovalifolia*. *Elench. H. R. M. 1805.*

Hab. in N. H. ☿. Semina communicavit. *D. Sord.*

### MELAMPEDIUM. Linn.

*Char. generis differ.*

*Anthodium* simplici serie pentaphyllum. *Corolla* radiata, ligulis ovatis, concavis, emarcescentibus, persistentibus. *Receptaculum* paleaceum. *Semina* calva, palea receptaculi induratis obvoluta.

*Obs. Palea receptaculi semina obvolventes apice in appendiculum squamiformem concavum pteromque declinat. Huc palearum appendiculum pro pappo pappum assumpere Linnæus, alique.*

404. *MELAMP. sericeum*: foliis subulatis, inferioribus trifidis, superioribus integris linearibus lanceolatis; seminum appendiculis retrorsum uncinatis.

*Melamp. sericeum*. *Elench. H. R. M. ann. 1805.*

Habitat in N. Hisp. V. V. in R. M. H. per plures annos, eandem ex seminibus à *prop. D. D. Sord et Melamp. sericeis*. ☿.

405. *MELAMP. heterophyllum*: caule erecto: foliis lanceolatis indivisis trifidisque subtus sericeis: seminibus muricatis inappendiculatis.

β. foliis omnibus indivisis.

Habit. in N. H. C.

#### ORD. 5. POLYGAMIA SEGREGATA.

406. *FLAVERIA repanda*: foliis ellipticis repando dentatis: caule subdiviso.

*Azumates* vulgo in Nova Hispania, ubi sponte nascitur.

Colitur in R. M. H. ubi Autumno floret. C.

407. *FLAVERIA linearis*: foliis linearibus iaregerantibus: ovario vaginantibus.

Genus *Boldo* herb.

*Subfrutex* spithameus, vel parum amplius, glaberrimus.

Venit ad maris litora prope *Havanam*, ubi *Boldo* legit.

*Obs.* Huc pertinent *Milleria contrayerba* et *angustifolia* cl. *Cav.*

#### CLASSIS XX. CRYPTOGRAMIA.

##### ORD. FILICES.

408. *ASPLENIUM leptophyllum*: frondibus lanceolatis bipinnatis: pinnatis dense approximatis, foliolis cuneiformibus 2-3-partitis, lobulo medio ovato, *Anal. de Class. Natur. vol. 5. pag. 155. tab. 41. fig. 3. optima.*

Habitat in Regno Valentino.

*Planta* genere, totoque caelo diversa ab *Hemionitide leptophylla*, cui perperam adjunxerunt *Clar. Swartz et Willden.*

409. *HEMIONITIS Pozoi*: frondibus pinnatis: pinnulis inferioribus sessilibus, distinctis: superioribus confluentibus.

Habitat in Cantabriae provinciis, ubi legit praelaudatus *D. F. del Pozo. l.*

410. *HEMIONITIS leptophylla*: frondibus pinnatis bipinnatisque glaberrimis, inferne nudis: pinnulis cuneiformibus rotundato-lobatis: soris discretis, et confluentibus.

*Polypodium leptophyllum*: *Linn. Sp. pl. Burr. rarior. tab. 431.*

*Grammitis leptophylla*. *Swart. Synops. Fil. pag. 23. et 118. tab. 1. f. 6. excluso Cav. Synonymo.*

Habit. locis saxosis in Hispania australiore, praesertim in Regno Cordubense.

G. N. LLAVEA.

##### Character generis.


FRUCTIFICATIO dorsalis in punctis oblongis, sive lineolis, nervo obliquis: initio indusio omnino tecta.

INDUSIUM marginale, membranaceum, continuum, interius dehiscens.

CAPSULAE pedicellatae, annulatae, uniloculares, bivalves: annulus elastice distiliens.

411. *LLAVEA cordifolia*. — *Frons* bipinnata: foliola sterilia, cordata, serrulata, fertilia linearia integerrima.

Habitat in Nova Hispania.







Parrote maverdis



COPIED QUARTER, QUARTER  
TA MARAVERDIS, ANO  
THE DOCHOCIENTES  
THE DOCHOC.

UNIVERSITY OF  
CALIFORNIA







S1826

14 DAY USE  
RETURN TO DESK FROM WHICH BORROWED

BIOLOGY LIBRARY

This book is due on the last date stamped below, or  
on the date to which renewed.

Renewed books are subject to immediate recall.

~~MAY 1 2 1965~~

~~INTER-LIBRARY  
LOAN~~

~~MAY 13 1965~~

~~MAY 7 1965~~

Due end of SPRING quarter  
Subject to recall after —

~~MAY 13 1974~~  
Due end of FALL quarter  
Subject to recall after —

~~SEP 24 1974~~

11-12-05

REC'D BIOS

SEP 12 '05 - 2:00 PM

LD 21-40m-4,'64  
(E4555s10)476

General Library  
University of California  
Berkeley



554734

QK92  
L3

UNIVERSITY OF CALIFORNIA LIBRARY

